arXiv:1006.5520v2 [math.CO] 26 Mar 2011

On duality and fractionality of multicommodity flows in directed networks

Hiroshi HIRAI
Department of Mathematical Informatics,
Graduate School of Information Science and Technology,
University of Tokyo, Tokyo, 113-8656, Japan
`hirai@mist.i.u-tokyo.ac.jp`

Shungo KOICHI
Department of Systems Design and Engineering,
Nanzan University, Seto 489-0863, Japan
`shungo@nanzan-u.ac.jp`

June 2010
March 2011 (revised)

Abstract

In this paper we address a topological approach to multiflow (multicommodity flow) problems in directed networks. Given a terminal weight μ, we define a metrized polyhedral complex, called the directed tight span T_μ, and prove that the dual of μ-weighted maximum multiflow problem reduces to a facility location problem on T_μ. Also, in case where the network is Eulerian, it further reduces to a facility location problem on the tropical polytope spanned by μ. By utilizing this duality, we establish the classifications of terminal weights admitting combinatorial min-max relation (i) for every network and (ii) for every Eulerian network. Our result includes Lomonosov-Frank theorem for directed free multiflows and Ibaraki-Karzanov-Nagamochi's directed multiflow locking theorem as special cases.

Keywords: multicommodity flows, metrics, min-max theorems, facility locations

1 Introduction

A *network* (G, S, c) is a triple of a directed graph $G = (VG, EG)$, a specified set $S \subseteq VG$ of nodes called *terminals*, and a nonnegative integer-valued edge-capacity $c : EG \to \mathbf{Z}_+$. An *S-path* is a (directed) path joining distinct terminals. A *multiflow* (*multicommodity flow*) is a pair $(\mathcal{P}, \lambda)$ of a set $\mathcal{P}$ of S-paths and a nonnegative flow-value function $\lambda : \mathcal{P} \to \mathbf{R}_+$ satisfying the capacity constraint: $\sum\{\lambda(P) \mid P \in \mathcal{P}, P \text{ contains } e\} \leq c(e)$ for $e \in EG$. Given a nonnegative terminal weight $\mu : S \times S \to \mathbf{R}_+$, the flow-value $\mathrm{val}(\mu, f)$ of multiflow $f = (\mathcal{P}, \lambda)$ is defined by $\sum\{\lambda(P)\mu(s_P, t_P) \mid P \in \mathcal{P}\}$, where s_P and t_P denote the start node and the end node of P, respectively. Then the *μ-weighted maximum multiflow problem* is formulated as:

μ-MFP: Maximize $\mathrm{val}(\mu, f)$ over all multiflows f in (G, S, c).

For a special terminal weight μ, the μ-MFP has a nice integrality property. For example, consider $S = \{s, t\}$ and $(\mu(s, t), \mu(t, s)) = (1, 0)$. Then the max-flow min-cut theorem says that the maximum flow value is equal to the minimum (s,t)-cut value and there always exists an *integral* maximum flow (maximum flow $(\mathcal{P}, \lambda)$ for which λ is integer-valued). Consider the case where $\mu(s,t) = 1$ for all distinct $s, t \in S$, and network is Eulerian. Lomonosov [14] and Frank [3] independently proved that the maximum flow value is equal to the sum of the minimum $(s, S \setminus s)$-cut value over $s \in S$ and there exists an integral maximum multiflow.

The goal of this paper is to classify weight functions $\mu : S \times S \to \mathbf{R}_+$ for which μ-MFP possesses such a combinatorial min-max relation. This classification problem, called the *fractionality problem*, was raised by Karzanov for the undirected μ-MFP (G is undirected and μ is symmetric); see [11]. It is well-known that the LP-dual to μ-MFP is a linear optimization over metrics on node set VG. In 90's, Karzanov [12, 13] found a remarkable fact that all possible candidates of optimal metrics are embedded into *a metric space on a polyhedral complex associated with* μ. This polyhedral complex is known as the *tight span*, which was earlier introduced by Isbell [10] and Dress [2] independently. Then the LP-dual reduces to a *facility location problem* on the tight span. Furthermore, if the tight span has a sufficiently nice geometry (dimension at most two), then one can obtain a combinatorial min-max relation from its shape. Otherwise (dimension at least three), one can conclude that μ-MFP has no such a combinatorial duality relation. Recently, this beautiful theory was further extended by the first author, and the fractionality problem for the undirected μ-MFP was roughly settled [5, 6, 7].

Our previous paper [8] started to develop an analogous duality theory for directed multiflows. In the directed case, the LP-dual is a linear optimization over *possibly asymmetric* metrics, which we call *directed metrics.* We introduced a directed version T_μ of the tight span (*directed tight span*). In the case of metric μ-MFP (μ is a directed metric), we showed that the LP-dual reduces to a facility location problem on T_μ; see [8, Section 4]. Moreover, in the case where a network is Eulerian, this LP-dual further reduces to a facility location problem on the *tropical polytope* $\bar{Q}_\mu$ spanned by μ, which was earlier introduced by Develin-Sturmfels [1] in the context of the tropical geometry.

The main contribution of this paper extends this duality theory for possibly nonmetric weights and solves the fractionality problems (i) for μ-MFP and (ii) for Eulerian μ-MFP (which is μ-MFP on an Eulerian network). In Section 2, we establish a general duality relation for μ-MFP with a possibly nonmetric weight μ. As well as the metric case, the LP-dual reduces to a facility location on the directed tight span T_μ (Theorem 2.2). However, in Eulerian case, we need a more careful treatment for the nonmetricity of μ. We newly introduce the *slimmed tropical polytope* $\bar{Q}_\mu^{slim}$, which is a certain subset of the tropical polytope and coincides with it if μ is a metric. Then we prove that the LP-dual to an Eulerian μ-MFP reduces to a facility location on $\bar{Q}_\mu^{slim}$ (Theorem 2.4). In Section 3, we show the integrality theorem (Theorem 3.1) that (i) if $\dim T_\mu \leq 1$, then every μ-MFP has an integral optimal multiflow, and (ii) if $\dim \bar{Q}_\mu^{slim} \leq 1$ then every Eulerian μ-MFP has an integral optimal multiflow. We remark that the former result can be proved by a reduction to the minimum cost circulation. The second result includes Lomonosov-Frank theorem for directed free multiflows [14, 3] and Ibaraki-Karzanov-Nagamochi's directed version of the multiflow locking theorem [9] as special cases. We give a combinatorial characterization of weights μ with $\dim \bar{Q}_\mu^{slim} \leq 1$ in terms of *oriented trees* (Theorem 3.4), and explain a relationship among these results. In Section 4, we show that the one-dimensionality of the directed tight span and the slimmed tropical polytope are best possible for the integrality. Theorem 4.1 says that if $\dim T_\mu \geq 2$, then there is no positive integer k such that every μ-MFP has a $1/k$-integral optimal multiflow, and that if $\dim \bar{Q}_\mu^{slim} \geq 2$, then there is no positive integer k such that every Eulerian μ-MFP has a $1/k$-integral optimal multiflow.

Notation. The sets of real numbers and nonnegative real numbers are denoted by $\mathbf{R}$ and $\mathbf{R}_+$, respectively. The set of functions from a set X to $\mathbf{R}$ (resp. $\mathbf{R}_+$) is denoted by $\mathbf{R}^X$ (resp. $\mathbf{R}^X_+$). For a subset $Y \subseteq X$, the characteristic function $\mathbf{1}_Y \in \mathbf{R}^X$ is defined by $\mathbf{1}_Y(x) = 1$ for $x \in Y$ and $\mathbf{1}_Y(x) = 0$ for $x \notin Y$. We particularly denote by $\mathbf{1}$ the all-one function in $\mathbf{R}^X$. For $p, q \in \mathbf{R}^X$, $p \leq q$ means $p(x) \leq q(x)$ for each $x \in X$, and $p < q$ means $p(x) < q(x)$ for each $x \in X$. For $p \in \mathbf{R}^X$, $(p)_+$ is defined by $((p)_+)(x) = \max\{p(x), 0\}$ for each $x \in X$. For a set P in $\mathbf{R}^X$, a point p in P is said to be *minimal* if there is no other point $q \in P \setminus p$ with $q \leq p$.

For a set S, a nonnegative real-valued function d on $S \times S$ having zero diagonals $d(s,s) = 0$ ($s \in S$) is called a *directed distance.* We regard a terminal weight $S \times S \to \mathbf{R}_+$ as a directed distance. A directed distance d on a set S is called a *directed metric* if it satisfies the triangle inequality $d(s,t) + d(t,u) \geq d(s,u)$ for every triple $s,t,u \in S$. A *directed metric space* is a pair (S,μ) of a set S and a directed metric d on S. For a directed metric d on V, and two subsets $A, B \subseteq V$, let $d(A,B)$ denote the minimum distance from A to B:

$$d(A,B) = \inf\{d(x,y) \mid (x,y) \in A \times B\}.$$

In our theory, the following directed metric D^+_∞ on $\mathbf{R}^X$ is particularly important:

$$D^+_\infty(p,q) = \|(q-p)_+\|_\infty \quad (= \max_{x \in X}(q(x) - p(x))_+) \quad (p, q \in \mathbf{R}^X).$$

We remark that $D^+_\infty(p,q) = 0$ whenever $p \geq q$.

For a directed or undirected graph G, its node set and edge set are denoted by VG and EG, respectively. If directed, an edge with tail x and head y is denoted by xy. If undirected, we do not distinguish xy and yx. In a network (G,S,c), a non-terminal node is called an *inner node.* For a node $x \in VG$, we say "*x fulfills the Eulerian condition*" if the sum of the capacities $c(xy)$ over edges xy leaving x is equal to that over edges entering x. A network (G,S,c) is said to be *inner Eulerian* if every inner node fulfills the Eulerian condition, and is said to be *totally Eulerian* if every node fulfills the Eulerian condition.

A directed distance and directed metric is often simply called a distance and a metric, respectively.

2 Duality

Let (G,S,c) be a network and let μ be a directed distance on S. We denote by $\mathrm{MFP}^*(\mu; G,S,c)$ the optimal value of μ-MFP for (G,S,c). The linear programing dual to μ-MFP is given by

$$\begin{array}{lll} \textbf{LPD}: & \text{Minimize} & \displaystyle\sum_{xy \in EG} c(xy)d(x,y) \\ & \text{subject to} & d \text{ is a directed metric on } VG, \\ & & d(s,t) \geq \mu(s,t) \quad (s,t \in S). \end{array}$$

We are going to represent LPD as a facility location problem on a metrized polyhedral complex associated with μ. Let S^c and S^r be copies of S. For an element $s \in S$, the corresponding elements in S^c and S^r are denoted by s^c and s^r, respectively. We denote $S^c \cup S^r$ by S^{cr}. For a point $p \in \mathbf{R}^{S^{cr}}$, the restrictions of p to S^c and S^r are denoted by p^c and p^r, respectively, i.e., $p = (p^c, p^r)$. Consider the following unbounded polyhedron in $\mathbf{R}^{S^{cr}}$:

$$P_\mu = \{p \in \mathbf{R}^{S^{cr}} \mid p(s^c) + p(t^r) \geq \mu(s,t) \ (s,t \in S)\}.$$

Let D_∞ be a directed metric on $\mathbf{R}^{S^{cr}}$ defined as

$$D_\infty(p,q) = \max\{D^+_\infty(p^c,q^c),\ D^+_\infty(q^r,p^r)\} \quad (p,q \in \mathbf{R}^{S^{cr}}).$$

We endow P_μ and its subsets with this directed metric. For a subset R in P_μ, we denote by $(R)^+ = R^+$ the set of nonnegative points in R. Also for $s \in S$ we denote by $(R)_s = R_s$ the set of points $p \in R$ with $p(s^c) + p(s^r) = \mu(s,s) = 0$; if $R \subseteq \mathbf{R}^{S^{cr}}_+$ then R_s is the set of points $p \in R$ with $p(s^c) = p(s^r) = 0$.

For a subset $R \subseteq P^+_\mu$, consider the following *facility location problem* on R:

$$\begin{array}{lll} \textbf{FLP}: & \text{Minimize} & \displaystyle\sum_{xy \in EG} c(xy) D_\infty(\rho(x),\rho(y)) \\ & \text{subject to} & \rho : VG \to R, \\ & & \rho(s) \in R_s \ (s \in S). \end{array}$$

Let $\mathrm{FLP}^*(R;G,S,c)$ denote the minimum value of this problem. Then the following weak/strong duality holds:

Lemma 2.1. *For a network (G,S,c) and a directed distance μ on S, we have*

(1) $\mathrm{MFP}^*(\mu;G,S,c) \leq \mathrm{FLP}^*(R;G,S,c)$ *for any subset* $R \subseteq P^+_\mu$*, and*

(2) $\mathrm{MFP}^*(\mu;G,S,c) = \mathrm{FLP}^*(P^+_\mu;G,S,c)$.

Proof. We first note the following property:

(2.1) For $s,t \in S$ and $(p,q) \in R_s \times R_t$ we have $D_\infty(p,q) \geq \mu(s,t)$.

Indeed, $D_\infty(p,q) \geq (q(s^c) - p(s^c))_+ \geq q(s^c) + q(t^r) \geq \mu(s,t)$, where we use $p(s^c) = q(t^r) = 0$.

It suffices to show (2). Take a map $\rho : VG \to P^+_\mu$ feasible to FLP. Let d be a metric on VG defined by $d(x,y) = D_\infty(\rho(x),\rho(y))$. By (2.1), we have $d(s,t) = D_\infty(\rho(s),\rho(t)) \geq \mu(s,t)$ for $s,t \in S$. Thus d is feasible to LPD with the same objective value. Conversely, take a metric d feasible to LPD. Define $\rho : VG \to \mathbf{R}^{S^{cr}}$ by $((\rho(x))(s^c),(\rho(x))(s^r)) = (d(s,x),d(x,s))$ for $s \in S$. Then $(\rho(x))(s^c) + (\rho(x))(t^r) = d(s,x) + d(x,t) \geq d(s,t) \geq \mu(s,t)$. Hence $\rho(x) \in P^+_\mu$. Moreover $(\rho(s))(s^c) = (\rho(s))(s^r) = d(s,s) = 0$. Thus ρ is feasible to FLP for $R = P^+_\mu$. By triangle inequality we have $D_\infty(\rho(x),\rho(y)) = \max\{\max_{s \in S}(d(s,y) - d(s,x))_+, \max_{t \in S}(d(x,t) - d(y,t))_+\} \leq d(x,y)$. Since c is nonnegative, we have $\sum c(xy) D_\infty(\rho(x),\rho(y)) \leq \sum c(xy) d(x,y)$. $\square$

In the following, we are going to determine "reasonably small" subsets $R \subseteq P^+_\mu$ for which the strong duality holds (i) for general networks and (ii) for Eulerian networks. In the next subsection (Section 2.1), we introduce the directed tight span T_μ and a fiber Q_μ of the tropical polytope as subsets in P_μ, and list their fundamental properties, shown by our previous paper [8]. In Section 2.2, we show that the strong duality holds for $R = T_\mu$ in every network (Theorem 2.2). We introduce a notion of a slimmed section and show that the strong duality holds for a slimmed section $R \subseteq Q_\mu$ in every Eulerian network (Theorem 2.4).

2.1 Preliminary: tight spans and tropical polytopes

Consider the following (non-convex) polyhedral subsets in P_μ:

$$\begin{array}{rcl} T_\mu & = & \text{the set of minimal elements of } P^+_\mu. \\ Q_\mu & = & \text{the set of minimal elements of } P_\mu. \end{array}$$


$(Q_\mu)_s$
Q_μ
$(Q_\mu)_u$
$(\mathbf{R}^2, D_\infty^+)$
Q_μ^+
R_s
$(Q_\mu)_t$
R
$\mathbf{R}^{S^{cr}}$
R_u
R_t
$(\mathbf{1}, -\mathbf{1})\mathbf{R}$
$\bar{Q}_\mu$
$\mathbf{R}^{S^{cr}}/(\mathbf{1}, -\mathbf{1})\mathbf{R}$

Figure 1: Q_μ, Q_μ^+, and $\bar{Q}_\mu$

We call T_μ the *directed tight span*. The polyhedron P_μ has the linearity space $(\mathbf{1}, -\mathbf{1})\mathbf{R}$. The projection $\bar{Q}_\mu := Q_\mu/(\mathbf{1}, -\mathbf{1})\mathbf{R}$ is known as the *tropical polytope* generated by matrix $(-\mu(s,t) \mid s,t \in S)$; see Develin and Sturmfels [1]. We note the relation: $Q_\mu \supseteq Q_\mu^+ \subseteq T_\mu \subseteq P_\mu$. In the inclusions, T_μ is a subcomplex (i.e., a union of faces) of P_μ, and Q_μ^+ is a subcomplex of T_μ. A subset $R \subseteq Q_\mu$ is called a *section* if the projection $p \in R \mapsto \bar{p} \in \bar{Q}_\mu$ is bijective. A subset $R \subseteq \mathbf{R}^{S^{cr}}$ is said to be *balanced* if there is no pair p, q of points in R such that $p^c < q^c$ or $p^r < q^r$. In fact, the projection $Q_\mu^+ \to \bar{Q}_\mu$ is surjective and there always exists a balanced section in Q_μ^+ [8, Lemma 2.4]. Figure 1 illustrates Q_μ, Q_μ^+, and $\bar{Q}_\mu$ for all-one distance on a 3-set $\{s,t,u\}$. In this case, $T_\mu = Q_\mu^+$ holds, Q_μ consists of three infinite strips with a common side, Q_μ^+ is a folder consisting of three triangles, and $\bar{Q}_\mu$ is a star of three leaves.

The rest of this subsection is devoted to listing basic properties of these polyhedral sets. They were proved in [8, Section 2]. The most important property for us is the existence of *nonexpansive retractions* among them.

A. Nonexpansive retractions. For two directed metric spaces (V,d) and (V',d'), a map $\phi : V \to V'$ is said to be *nonexpansive* if $d'(\phi(x), \phi(y)) \leq d(x,y)$ for all pairs $x, y \in V$. By a *cycle* C of V we mean a cyclic permutation $(x_1, x_2, \ldots, x_n)$ of a finite (multi-)set in V. Its *length* $d(C)$ is defined by $d(x_1,x_2) + d(x_2,x_3) + \cdots + d(x_{n-1},x_n) + d(x_n,x_1)$. Also, $\phi : V \to V'$ is said to be *cyclically nonexpansive* if $d'(\phi(C)) \leq d(C)$ for all cycles C in V. A map from a set V to its subset $S \subseteq V$ is said to be a *retraction* if it is identity on S.

(2.2) (1) There exists a nonexpansive retraction $\phi : P_\mu^+ \to T_\mu$ with $\phi(p) \leq p$ for $p \in P_\mu^+$.

(2) There exists a cyclically nonexpansive retraction $\varphi : T_\mu \to Q_\mu^+$.

(3) For any balanced section $R \subseteq Q_\mu$, the retraction $\varphi_R : Q_\mu \to R$ determined by the relation

$$\varphi_R(p) - p \in (\mathbf{1}, -\mathbf{1})\mathbf{R} \quad (p \in Q_\mu)$$

is cyclically nonexpansive.

See Figure 2 for the retraction in (3).

B. Geodesics and embedding. A path $P \subseteq \mathbf{R}^{S^{cr}}$ is the image of a continuous map $\varrho : [0,1] \to \mathbf{R}^{S^{cr}}$. The length of P from $\varrho(0)$ to $\varrho(1)$ is defined by the supremum of $\sum_{i=0}^{n-1} D_\infty(\varrho(t_i), \varrho(t_{i+1}))$ over all $n > 0$ and $0 = t_0 < t_1 < \cdots < t_n = 1$. For simplicity, we restrict ϱ to be sufficiently nice: ϱ is injective and its length is finite. A subset $R \subseteq \mathbf{R}^{S^{cr}}$ is said to be *geodesic* if each pair $p, q \in R$ of points is joined by a path in R having length $D_\infty(p, q)$ from p to q.

(2.3) T_μ, Q_μ, Q_μ^+ and any balanced section in Q_μ are all geodesic.

For $s \in S$, let μ_s be the point in $\mathbf{R}^{S^{cr}}$ defined by

$$(\mu_s(t^c), \mu_s(t^r)) = (\mu(t,s), \mu(s,t)) \quad (t \in S). \tag{2.4}$$

Namely μ_s is composed by s-th column and s-th row vectors of μ (as a matrix).

(2.5) (1) For any balanced section R in Q_μ^+ we have $R_s = (Q_\mu)_s^+ = (T_\mu)_s$.

(2) For $s, t \in S$ we have $D_\infty((T_\mu)_s, (T_\mu)_t) = \mu(s,t)$.

(3) If μ is a metric, then $(T_\mu)_s = \{\mu_s\}$ for each $s \in S$.

In particular, if μ is a metric, then metric space (S, μ) is isometrically embedded into any balanced section R in Q_μ^+ by $s \mapsto \mu_s$.

C. Tight extensions. For a metric μ on S, an *extension* of μ is a metric d on V with $S \subseteq V$ and $d(s,t) = \mu(s,t)$ for $s, t \in S$, and it is said to be *tight* if there is no other extension d' on V with $d' \neq d$ and $d' \leq d$. Also an extension d on V of μ is said to be *cyclically tight* if there is no other extension d' on V such that $d'(C) \leq d(C)$ for all cycles C in V and $d'(C) < d(C)$ for some cycle C. Every cyclically tight extension is a tight extension. The converse is not true. For example, $S = \{s, t\}$, $\mu(s,t) = \mu(t,s) = 1$, $V = \{s, t, u, v\}$, and consider two extensions d, d' on V defined by

$$d = \begin{array}{|c|cccc|} \hline & s & t & u & v \\ \hline s & 0 & 1 & 1 & 0 \\ t & 1 & 0 & 1 & 0 \\ u & 0 & 0 & 0 & 0 \\ v & 1 & 1 & 1 & 0 \\ \hline \end{array}, \qquad d' = \begin{array}{|c|cccc|} \hline & s & t & u & v \\ \hline s & 0 & 1 & 1 & 1 \\ t & 1 & 0 & 1 & 1 \\ u & 0 & 0 & 0 & 0 \\ v & 0 & 0 & 0 & 0 \\ \hline \end{array}$$

Then d is tight, but not cyclically tight. Indeed, for every pair $(x, y) \in V \times V \setminus S \times S$ if $d(x,y) > 0$, then $d(s,t) = d(s,x) + d(x,y) + d(y,t)$ or $d(t,s) = d(t,x) + d(x,y) + d(y,s)$; this means that we cannot decrease $d(x,y)$ keeping the triangle inequality. Compare d

with d'. Then $d'(C) \leq d(C)$ for all cycles C, and $0 = d'(u,v) + d'(v,u) < d(u,v) + d(v,u) = 1$. Thus d is not cyclically tight.

Every tight extension and cyclically tight extension are embedded into (T_μ, D_∞) and (Q_μ^+, D_∞), respectively. We use this fact in Section 4.

(2.6) Let μ be a metric on S and d its extension on V.

(1) d is tight if and only if there is an isometric embedding $\rho : V \to T_\mu$ such that $\rho(s) = \mu_s$ for each $s \in S$.

(2) d is cyclically tight if and only if there is an isometric embedding $\rho : V \to Q_\mu^+$ such that $\rho(s) = \mu_s$ for each $s \in S$ and $\rho(V)$ is balanced.

Here an *isometric embedding* from (V,d) to (V',d') is a map $\rho : V \to V'$ satisfying $d'(\rho(x), \rho(y)) = d(x,y)$ for all $x, y \in V$.

D. Further technical stuffs. For a point $p \in P_\mu$, let $K_\mu(p) = K(p)$ denote the bipartite graph on S^{cr} with edge set $\{s^c t^r \mid p(s^c) + p(t^r) = \mu(s,t)\}$.

(2.7) (1) A point $p \in P_\mu^+$ belongs to T_μ if and only if $K(p)$ has no isolated node u with $p(u) > 0$.

(1') A point $p \in P_\mu$ belongs to Q_μ if and only if $K(p)$ has no isolated node.

(2) For $p \in T_\mu$, the dimension of the minimal face of T_μ containing p is equal to the number of components in $K(p)$ having no node u with $p(u) = 0$.

(2') For $p \in Q_\mu$, the dimension of the minimal face of Q_μ containing p is equal to the number of components in $K(p)$ [1, Proposition 17].

(3) Any k-dimensional face F in (T_μ, D_∞) is isometric to a k-dimensional polytope in $(\mathbf{R}^k, D_\infty^+)$.

(4) $D_\infty(p,q) = D_\infty^+(p^c, q^c) = D_\infty^+(q^r, p^r)$ holds for $p, q \in Q_\mu$ and for $p, q \in T_\mu$.

(5) T_μ has dimension at most 1 if and only if T_μ is a path isometric to a segment in $(\mathbf{R}, D_\infty^+)$.

The property (3) follows from [8, (2.1)].

Our technical arguments use a method of perturbing a point $p \in Q_\mu^+$ to another point $p' \in Q_\mu^+$. For a node subset U in a graph $K(p)$, the set of nodes in $S^{cr} \setminus U$ incident to U is denoted by $N_p(U) = N(U)$. The following consideration is a basis for our perturbation method, which has a similar flavor of manipulating dual variables in bipartite matching problems:

(2.8) For $p \in Q_\mu^+, X \subseteq S$, let $p' := p - \epsilon \mathbf{1}_{X^c}$ for small $\epsilon > 0$, and $Y^r := N_p(X^c)$.

- $p(s^c) > 0$ ($s^c \in X^c$) is necessary for keeping the nonnegativity of p'.
- Put $p' \leftarrow p' + \epsilon \mathbf{1}_{Y^r}$ so that $p' \in P_\mu$.
- Then all edges joining $S^c \setminus X^c$ and Y^r vanish in $K(p')$.
- Therefore, p' belongs to Q_μ^+ if and only if each node in $S^c \setminus X^c$ is joined to $S^r \setminus Y^r$ in $K(p)$.
- We can increase ϵ until some coordinate of p' in X^c reaches zero or there appears an edge joining X^c and $S^r \setminus Y^r$.

Here the fourth implication uses (2.7) (1').

2.2 Duality relations

First we establish strong duality relations for general networks and for inner Eulerian networks, which are easy consequences of the existence of nonexpansive retractions (2.2).

Theorem 2.2. *Let μ be a directed distance on S.*

(1) $\mathrm{MFP}^*(\mu; G, S, c) = \mathrm{FLP}^*(T_\mu; G, S, c)$ *holds for every network* (G, S, c).

(2) $\mathrm{MFP}^*(\mu; G, S, c) = \mathrm{FLP}^*(R; G, S, c)$ *holds for every balanced section R in Q_μ^+ and every inner Eulerian network* (G, S, c).

Proof. Take an optimal map ρ for FLP with $R = P_\mu^+$. Take a nonexpansive retraction $\phi : P_\mu^+ \to T_\mu$ in (2.2) (1). Consider the composition $\phi \circ \rho : VG \to T_\mu$. Then $\phi \circ \rho$ is also feasible, and does not increase the objective value. Thus we have (1).

Next we show (2). Suppose that (G, S, c) is inner Eulerian. Then the capacity function $c : EG \to \mathbf{Z}_+$ is decomposed into the sum of the incidence vectors of cycles $C_1, C_2, \ldots, C_m$ and S-paths $P_1, P_2, \ldots, P_n$ (possibly repeating). Take an optimal map ρ for FLP with $R = T_\mu$. Then we have

$$\sum_{xy \in EG} c(xy) D_\infty(\rho(x), \rho(y)) = \sum_{i=1}^{m} D_\infty(\rho(C_i)) + \sum_{j=1}^{n} D_\infty(\rho(P_j)). \tag{2.9}$$

We can take a cyclically nonexpansive retraction $\varphi : T_\mu \to R$ by (2),(3) in (2.2). Since $\varphi \circ \rho(s) = \rho(s) \in (T_\mu)_s = R_s$ for each $s \in S$ by (2.5) (1), φ is identity on $(T_\mu)_s$ and thus $\varphi \circ \rho$ is also feasible. Moreover, by cyclically nonexpansiveness, $D_\infty(\varphi \circ \rho(C_i)) \leq D_\infty(\rho(C_i))$ holds and also $D_\infty(\varphi \circ \rho(P_j)) \leq D_\infty(\rho(P_j))$ holds by $D_\infty(\varphi \circ \rho(s), \varphi \circ \rho(t)) = D_\infty(\rho(s), \rho(t))$ for $s, t \in S$. Hence $\varphi \circ \rho$ is also optimal. □

We give some examples. Consider the all-one distance μ on a 3-set $\{s, t, u\}$; recall Figure 1. Then the FLP is a location problem on a directed metric space on a folder consisting of three triangles, each of which is isometric to triangle $\{(x, y) \in \mathbf{R}^2 \mid 0 \leq y \leq x \leq 1\}$ in $(\mathbf{R}^2, D_\infty^+)$. Suppose that the network is inner Eulerian. We can take a balanced section R in Q_μ^+, which is a tree. By a cyclically nonexpansive retraction from Q_μ^+ to R, the FLP reduces to a location problem on the tree R; see Figure 2. Consider the 2-commodity flow case; let $S = \{s, s', t, t'\}$, and let $\mu(s, t) = \mu(s', t') = 1$ and let the other distances be zero. Then T_μ is given by $\{\mathbf{1}_{\{t,t'\}^r} + \alpha(\mathbf{1}_{s^c} - \mathbf{1}_{t^r}) + \beta(\mathbf{1}_{(s')^c} - \mathbf{1}_{(t')^r}) \mid 0 \leq \alpha, \beta \leq 1\}$, which is isometric to a square $\{(x, y) \in \mathbf{R}^2 \mid 0 \leq x, y \leq 1\}$ in $(\mathbf{R}^2, D_\infty^+)$.



Figure 2: Retraction from Q_μ^+ to R



Figure 3: The tight span T_μ for the 2-commodity distance μ

Terminal regions $(T_\mu)_s, (T_\mu)_t, (T_\mu)_{s'}, (T_\mu)_{t'}$ correspond to four sides as in Figure 3. In this case, $T_\mu = Q_\mu^+$ holds, and moreover Q_μ^+ itself is a balanced section. In contrast to the previous example, the region contraction in FLP does not occur if the inner Eulerian condition is imposed.

Slimmed sections and Eulerian condition on terminals. Next we consider the case where some of terminals fulfill the Eulerian condition. In this case, the strong duality holds for further smaller subsets in Q_μ, called *slimmed sections*. To define a slimmed section, we need several (somewhat technical) notions. Recall notions of $K(p)$ and $N_p(\cdot)$ associated with $p \in Q_\mu$; see Section 2.1 D. Let $\mathcal{S}_0$ be the set of subsets X of S such that $\mu(s,t) = 0$ for all $(s,t) \in X \times X$; obviously $\{s\} \in \mathcal{S}_0$. For $X \in \mathcal{S}_0$, let $Q_{\mu,X}$ denote the set of points $p \in Q_\mu$ with $s^c s^r \in EK(p)$ for $s \in X$ and $s^c s^r \notin EK(p)$ for $s \notin X$; in particular $Q_{\mu,X} = \bigcap_{s \in X} (Q_\mu)_s \setminus \bigcup_{s \in S \setminus X} (Q_\mu)_s$. A point p in Q_μ is called a *fat* relative to X if $p \in Q_{\mu,X}$, and $N_p(S^c \setminus X^c) \subseteq S^r \setminus X^r$ or $N_p(S^r \setminus X^r) \subseteq S^c \setminus X^c$. The *degenerate set* $Q_{\mu,X}^{deg}$ relative to X is the set of points p in $Q_{\mu,X}$ with $N_p(S^c \setminus X^c) = S^r \setminus X^r$ or $N_p(S^r \setminus X^r) = S^c \setminus X^c$. Any point in a degenerate set is a fat. A *proper fat* is a fat not belonging to any degenerate set. Let Q_μ^{slim} be the subset of Q_μ obtained by deleting all proper fats. We consider the following equivalence relation $\sim$ on Q_μ^{slim}: $p \sim q$ if $p - q \in (\mathbf{1}, -\mathbf{1})\mathbf{R}$, or for some $X \in \mathcal{S}_0$, both p and q belong to $Q_{\mu,X}^{deg}$ and $p - q \in (\mathbf{1}, -\mathbf{1})\mathbf{R} + (\mathbf{1}_{X^c}, -\mathbf{1}_{X^r})\mathbf{R}$. The quotient $Q_\mu^{slim}/\sim$ is called the *slimmed tropical polytope* associated with μ, and is denoted by $\bar{Q}_\mu^{slim}$. The tropical polytope and the slimmed tropical polytope are the same if μ is a metric.

Proposition 2.3. *If μ is a metric, then Q_μ has no fat, and hence $\bar{Q}_\mu^{slim} = \bar{Q}_\mu$.*



$$\mu = \begin{array}{c|cccc} & s & t & u & v \\ \hline s & 0 & 1 & 1 & 1 \\ t & 1 & 0 & 1 & 0 \\ u & 1 & 1 & 0 & 0 \\ v & 0 & 0 & 0 & 0 \end{array} \qquad \mu = \begin{array}{c|ccc} & s & t & u \\ \hline s & 0 & 2 & 1 \\ t & 1 & 0 & 0 \\ u & 0 & 0 & 0 \end{array}$$

Figure 4: Two examples of Q_μ^+

Proof. By $(Q_\mu)_s \supseteq Q_{\mu,X}$ for $s \in X$ and (2.5) (3), $Q_{\mu,X}^+$ is a single point μ_s for $s \in X$. In $K(\mu_s)$, node s^c is incident to all nodes in S^r, and s^r is incident to all nodes in S^c; thus μ_s is never a fat. $\square$

Again we consider special sections in Q_μ^{slim}, called *slimmed sections.* Here a section is a subset of Q_μ^{slim} bijectively projected into $\bar{Q}_\mu^{slim}$. We first define a slimmed section R in $(Q_\mu^{slim})^+$, which is a section such that it is balanced, and for each $X \in \mathcal{S}_0$, there is no pair $p, q \in (Q_{\mu,X}^{deg})^+$ such that $p(s^c) < q(s^c)$ for all $s^c \in S^c \setminus X^c$ or $p(s^r) < q(s^r)$ for all $s^r \in S^r \setminus X^r$. Next, a slimmed section R in Q_μ^{slim} is a section such that it is balanced and $R/(\mathbf{1}, -\mathbf{1})\mathbf{R} = R'/(\mathbf{1}, -\mathbf{1})\mathbf{R}$ for some slimmed section R' in $(Q_\mu^{slim})^+$ (recall that the projection from Q_μ^+ to $\bar{Q}_\mu$ is surjective).

Figure 4 depicts two examples of Q_μ^+ together with $K(p)$ for an interior point p in each face. In the left example, Q_μ^+ is obtained from a folder of two triangles by attaching one segment on the top. Here any point in triangles except upper edges is a proper fat relative to $\{v\}$. So $(Q_\mu^{slim})^+$ is a star of three edges, and is a slimmed section. In the right example, Q_μ^+ is the union of square and segment. Although there is no proper fat, points in the square except the left and right corners form a section of degenerate set $Q_{\mu,\{u\}}^{deg}$. A slimmed section is obtained by replacing the square by an appropriate curve connecting the left and right corners; see Figure 5.

A terminal $s \in S$ is said to be *proper* if $(Q_\mu)_s$ has no fat. A network (G, S, c) is said to be *properly inner Eulerian* (relative to μ) if every node except proper terminals fulfills the Eulerian condition. The main result here is the following:

Theorem 2.4. *Let μ be a directed distance on S.*

(1) $\mathrm{MFP}^*(\mu; G, S, c) = \mathrm{FLP}^*(R; G, S, c)$ *holds for every slimmed section R in $(Q_\mu^{slim})^+$ and every properly inner Eulerian network (G, S, c).*

(2) $\mathrm{MFP}^*(\mu; G, S, c) = \mathrm{FLP}^*(R; G, S, c)$ *holds for every slimmed section R in Q_μ^{slim} and every totally Eulerian network (G, S, c).*



Figure 5: Retraction from Q_μ^+ to a slimmed section

The proof uses the following new retraction lemma:

Lemma 2.5. *For any slimmed section R in Q_μ^+, there exists a cyclically nonexpansive retraction φ from Q_μ^+ to R with $\varphi((Q_\mu)_s^+) \subseteq (Q_\mu)_s^+$ for each $s \in S$.*

The proof of this lemma is given in the end of this subsection. Assuming Lemma 2.5, we prove Theorem 2.4. Let S^* be the set of proper terminals. Take an optimal map $\rho : VG \to Q_\mu^+$ for FLP. As in the proof of Theorem 2.2, there are cycles C_i and S^*-paths P_j such that (2.9) holds. Take a cyclically nonexpansive retraction φ in Lemma 2.5. Then $\varphi \circ \rho : VG \to R$ is feasible to FLP with R. Since φ is cyclically nonexpansive, $D_\infty(\varphi \circ \rho(C_i)) \leq D_\infty(\rho(C_i))$ and $D_\infty(\varphi \circ \rho(P_j)) \leq D_\infty(\rho(P_j))$, where the second inequality follows from the fact that φ is identity on R_s for each proper terminal $s \in S^*$. Thus $\varphi \circ \rho$ and ρ have the same objective value. The statement (2) follows from (1) and (2.2) (3).

Figure 5 illustrates cyclically nonexpansive retractions in the examples of Figure 4. Again the 2-commodity tight span in Figure 3 has no fat; the region contraction in FLP does not occur even if the totally Eulerian condition is imposed.

As a corollary, we obtain topological properties of slimmed sections:

Corollary 2.6. *Let $R \subseteq Q_\mu$ be a slimmed section.*

(1) *R is contractible and geodesic, and so is R_s for $s \in S$.*

(2) *If $R \subseteq Q_\mu^+$, then $\mu(s,t) = D_\infty(R_s, R_t)$ for $s, t \in S$.*

Proof. (1). A cyclically nonexpansive map is continuous in the Euclidean topology [8, Remark 2.4]. So R is homotopy equivalent to convex set P_μ^+, which is contractible. Since P_μ^+ is geodesic, so is R; see [8, Section 2.3]. Since R_s is a retract of a face of P_μ^+, it is contractible and geodesic by the same argument.

(2). Consider the Eulerian network (G, S, c) such that $c(st) = c(ts) = 1$ and the other capacities are zero. Obviously $\mathrm{MFP}^*(G, S, c) = \mu(s,t) + \mu(t,s)$. By $\mathrm{MFP}^*(\mu; G, S, c) = \mathrm{FLP}^*(R; G, S, c)$, there is $(p,q) \in R_s \times R_t$ with $D_\infty(p,q) + D_\infty(q,p) = \mu(s,t) + \mu(t,s)$. Necessarily $D_\infty(p,q) = \mu(s,t)$ and $D_\infty(q,p) = \mu(t,s)$ by (2.1). ☐

Proof of Lemma 2.5. In the proof, we denote by ${Q_\mu}^c$ and ${Q_\mu}^r$ the projections of Q_μ to $\mathbf{R}^{S^c}$ and $\mathbf{R}^{S^r}$, respectively. These projections are bijective and isometric by (2.7) (4). For $q \in {Q_\mu}^c$, we can lift q to $p \in Q_\mu$ with $p^c = q$ by $p(t^r) = \max_{s^c \in S^c}(\mu(s,t) - q(s^c))$.

We remark that $p(s^r) = p(s^c) = 0$ for $p \in Q_{\mu,X}^+$ and $s \in X \in \mathcal{S}_0$. Let B be any balanced section in Q_μ^+. By $Q_{\mu,X} \subseteq \bigcap_{s \in X}(Q_\mu)_s$ and (2.5) (1), B includes $Q_{\mu,X}^+$ for all $X \in \mathcal{S}_0$. One can verify from (2.8) that a point $p \in Q_{\mu,X}^+$ is a fat if and only if for small

$\epsilon > 0$ we have $p^c - \epsilon \mathbf{1}_{(S\setminus X)^c} \in (Q^+_{\mu,X})^c$ or $p^r - \epsilon \mathbf{1}_{(S\setminus X)^r} \in (Q^+_{\mu,X})^r$ (use the property that any node u with $p(u) = 0$ is incident to X^{cr} by nonnegativity of μ).

Based on this, for $\epsilon \geq 0$, consider the map $\varphi^c_{X,\epsilon}$ on B obtained by the following process. For each $p \in Q^+_{\mu,X}$, add $-\epsilon^* \mathbf{1}_{(S\setminus X)^c}$ to p^c, where ϵ^* is the maximum nonnegative real in $[0,\epsilon]$ such that $p^c - \epsilon^* \mathbf{1}_{(S\setminus X)^c}$ belongs to the closure of $(Q^+_{\mu,X})^c$. Then lift the resulting point q to $p' \in Q^+_{\mu,X}$ with $(p')^c = q$, and define $\varphi^c_{X,\epsilon}(p) := p'$. Extend $\varphi^c_{X,\epsilon}$ to a map $B \to B$ by defining it to be identity on the points not in $Q^+_{\mu,X}$.

In above p, ϵ, ϵ^*, the following key property holds:

(2.10) For $q \in B$ with $q = \varphi^c_{X,\epsilon}(q)$, if q is not a proper fat relative to any proper subset $Y \subset X$, then we have

$$\begin{array}{rcl} D_\infty(\varphi^c_{X,\epsilon}(p,q)) & \leq & D_\infty(p,q) + \epsilon^*, \\ D_\infty(\varphi^c_{X,\epsilon}(q,p)) & = & D_\infty(q,p) - \epsilon^*. \end{array}$$

Proof. By (2.7) (4), we may consider p^c, q^c, D^+_∞ instead of p, q, D_∞. The first relation is obvious from definition of D^+_∞; see [8, p. 8]. We show the second. We claim that $K(q)$ has an edge $s^c t^r$ joining $S^c \setminus X^c$ and X^r. Suppose not. Then $X^c \supseteq N_q(X^r)$. Since $X \in \mathcal{S}_0$, $\mu(s,t) = 0$ holds for all $(s^c, t^r) \in N_q(X^r) \times X^r$. Hence we have $q(u) = 0$ for $u \in N_q(X^r) \cup X^r$ and $q(s^c) > 0$ for $s^c \in S^c \setminus X^c$. Since $q = \varphi^c_{X,\epsilon}(q)$, necessarily $X^c \supset N_q(X^r)$ (proper inclusion), and q is a proper fat relative to Y with $Y^c = N_q(X^r) \subset X^c$; a contradiction.

For an edge $s^c t^r \in EK(q)$ joining $S^c \setminus X^c$ and X^r, we have $\epsilon^* \leq p(s^c) + p(t^r) - \mu(s,t) = p(s^c) + p(t^r) - q(s^c) - q(t^r) = p(s^c) - q(s^c) - q(t^r)$, where we use $q(s^c) + q(t^r) = \mu(s,t)$ by $s^c t^r \in EK(q)$ and $p(t^r) = 0$. Therefore we have $(*)$ $p(s^c) - q(s^c) \geq \epsilon^*$. Thus we have

$$\begin{array}{rcl} D^+_\infty(q^c, p^c - \epsilon^* \mathbf{1}_{(S\setminus X)^c}) & = & \|(p^c - \epsilon^* \mathbf{1}_{(S\setminus X)^c} - q^c)_+\|_\infty \\ & = & \max\limits_{t^c \in (S\setminus X)^c} \{(p(t^c) - \epsilon^* - q(t^c))_+\} = D^+_\infty(q^c, p^c) - \epsilon^*, \end{array}$$

where the second equality uses $p(t^c) = 0$ for all $t^c \in X^c$ and the third uses $(*)$. $\square$

We can define $\varphi^r_{X,\epsilon} : B \to B$ by changing roles of c and r, and an analogous property holds. Let $\varphi^c_X := \lim_{\epsilon \to \infty} \varphi^c_{X,\epsilon}$ and $\varphi^r_X := \lim_{\epsilon \to \infty} \varphi^r_{X,\epsilon}$ (well-defined). Next we study the image of φ^c_X. Let $p^* := \varphi^c_X(p)$ for $p \in Q^+_{\mu,X}$. Then $K(p^*)$ necessarily has an edge joining $S^c \setminus X^c$ and X^r. Therefore if p^* is a fat, then it is a fat relative to $Y \supset X$ (proper inclusion), or p^* is *r-maximal* in $(Q^{deg}_{\mu,X})^+$ in the sense that $p^* - \epsilon(\mathbf{1}_{(S\setminus X)^c}, -\mathbf{1}_{(S\setminus X)^r}) \notin (Q^{deg}_{\mu,X})^+$ for every $\epsilon > 0$. Also if p belongs to $(Q^{deg}_{\mu,X})^+$, then p^* is an r-maximal point with $p - p^* \in (\mathbf{1}_{(S\setminus X)^c}, -\mathbf{1}_{(S\setminus X)^r})\mathbf{R}$; see the right of Figure 5. Again an analogous property holds for φ^r_X by changing roles of r and c. Let $\varphi_X := \varphi^r_X \circ \varphi^c_X$. Then the image $\varphi_X(B)$ does not contain a proper fat relative to X.

Let B^{slim}_c be the subset of B obtained by deleting all proper fats and replacing each $(Q^{deg}_{\mu,X})^+$ by the set of its c-maximal points. Order all subsets $X_1, X_2, \ldots, X_m$ in $\mathcal{S}_0$ so that $X_i \subseteq X_j$ implies $i \leq j$. Then the composition $\varphi := \varphi_{X_m} \circ \varphi_{X_{m-1}} \circ \cdots \circ \varphi_{X_1}$ is a retraction from B to B^{slim}_c.

We show that $\varphi : B \to B^{slim}_c$ is cyclically nonexpansive. Let $X := X_i$. Take a cycle C in $\varphi_{X_{i-1}} \circ \varphi_{X_{i-2}} \circ \cdots \circ \varphi_{X_1}(B)$. We prove that $g(\epsilon) := D_\infty(\varphi^c_{X,\epsilon}(C)) - D_\infty(C)$ is a monotone nonincreasing function. It suffices to show $g(\epsilon) \leq 0$ for small $\epsilon > 0$. By construction, C does not contain proper fats relative to any $Y \subset X$. By (2.10), for a consecutive pair (p, q) in C we have

$$(2.11) \qquad D_\infty(\varphi^c_{X,\epsilon}(p, q)) - D_\infty(p, q) \begin{cases} \leq \epsilon & \text{if } p \neq \varphi^c_X(p), q = \varphi^c_X(q), \\ = -\epsilon & \text{if } p = \varphi^c_X(p), q \neq \varphi^c_X(q), \\ = 0 & \text{otherwise.} \end{cases}$$

Since the number of consecutive pairs (p, q) with $p \neq \varphi^c_X(p), q = \varphi^c_X(q)$ is equal to that with $p = \varphi^c_X(p), q \neq \varphi^c_X(q)$, summing up (2.11) over all consecutive pairs yields $g(\epsilon) \leq 0$. The argument for φ^r_X is similar. Thus φ is cyclically nonexpansive.

We next verify that B^{slim}_c is slimmed. Indeed, take an arbitrary pair p, p' of c-maximal points in $(Q^{deg}_{\mu,X})^+$. Then there are edges $s^c t^r \in EK(p), \tilde{s}^c \tilde{t}^r \in EK(p')$ joining X^c and $S^r \setminus X^r$. Hence $p'(s^c) + p'(t^r) = p'(t^r) \geq \mu(s, t) = p(t^r)$, and $p(\tilde{t}^r) \geq \mu(\tilde{s}, \tilde{t}) = p'(\tilde{t}^r)$.

Finally we construct a cyclically nonexpansive retraction from Q^+_μ to any slimmed section. Any slimmed section R in $(Q^{slim}_\mu)^+$ is obtained from a balanced section B by deleting all proper fats and replacing each $(Q^{deg}_{\mu,X})^+$ by a subset R_X with the properties that (i) there is no pair $p, q \in R_X$ with $p(s^c) < q(s^c)$ for all $s^c \in S^c \setminus X^c$ or $p(s^r) < q(s^r)$ for all $s^r \in S^r \setminus X^r$, and (ii) for each $p' \in (Q^{deg}_{\mu,X})^+$ there uniquely exists $p \in R_X$ with $p - p' \in (\mathbf{1}_{(S\setminus X)^c}, -\mathbf{1}_{(S\setminus X)^r})\mathbf{R}$. It suffices to give a cyclically nonexpansive map from B^{slim}_c to R. For each $X \in \mathcal{S}_0$, we can define a map φ^X_R on B^{slim}_c as: For each $p \in (Q^{deg}_{\mu,X})^+$, define $\varphi^X_R(p)$ to be the point p' in R_X determined by the relation $p' - p \in (\mathbf{1}_{(S\setminus X)^c}, -\mathbf{1}_{(S\setminus X)^r})\mathbf{R}$, and to be identity on the other points. So it suffices to prove that φ^X_R is cyclically nonexpansive; consider the composition of φ^X_R for all $X \in \mathcal{S}_0$. One can verify this fact in the essentially same way as above. The projection of $(Q^{deg}_{\mu,X})^+$ to $\mathbf{R}^{(S\setminus X)^{cr}}$ is isometry, and the image of R_X is a balanced set in $\mathbf{R}^{(S\setminus X)^{cr}}$. So we can apply the method in the proof of [8, Lemma 2.7]; the details are left to readers.

3 Integrality

The geometry of T_μ and $\bar{Q}^{slim}_\mu$ crucially affects the integrality of μ-MFP. The dimension of T_μ is defined by the largest dimension of faces of T_μ. The dimension of $\bar{Q}^{slim}_\mu$ is defined by the largest dimension of faces F of Q^{slim}_μ in modulo $\sim$; intuitively, it is the dimension of its slimmed section. The main goal of this section is to prove the following integrality theorem:

Theorem 3.1. *Let μ be a directed distance on S.*

1. *If $\dim T_\mu \leq 1$, then μ-MFP has an integral optimal multiflow for every network (G, S, c).*
2. *If $\dim \bar{Q}^{slim}_\mu \leq 1$, then μ-MFP has an integral optimal multiflow for every properly inner Eulerian network (G, S, c) (relative to μ).*

The first statement (1) is reducible to the minimum cost circulation. So we mainly concentrate on the second statement (2) and its consequences. The rest of this section is organized as follows. In next Section 3.1, we give basic definitions for cuts, cut distances, and oriented-tree realizations. Then, in Section 3.2, we prove Theorem 3.1 (2). In Section 3.3, we give a useful "combinatorial version" of Theorem 3.1 (2), and derive (slight) extensions of Lomonosov-Frank theorem for directed free multiflows and Ibaraki-Karzanov-Nagamochi's directed version of the multiflow locking theorem. In Section 3.4, we prove (1) by a reduction to the minimum cost circulation.

3.1 Preliminary: partial cuts, cut distances, and oriented trees

A *partial cut* on a set S is an ordered pair (A, B) of disjoint subsets $A, B \subseteq S$. We particularly call (A, B) a *cut* if $A \cup B = S$. For a partial cut (A, B) on S, the *cut distance* $\delta_{A,B} : S \times S \to \mathbf{R}_+$ is defined by

$$\delta_{A,B}(s,t) = \begin{cases} 1 & \text{if } (s,t) \in A \times B, \\ 0 & \text{otherwise,} \end{cases} \qquad (s, t \in S).$$

In a network (G, S, c), for a node subset $X \subseteq VG$, let ∂X denote the set of edges leaving X. For a partial cut (A, B) on S, the following relation is nothing but the max-flow min-cut theorem:

$$\text{MFP}^*(\delta_{A,B}; G, S, c) = \min\{c(\partial X) \mid A \subseteq X \subseteq VG \setminus B\}. \tag{3.1}$$

An *oriented tree* Γ is a directed graph whose underlying undirected graph is a tree. For a nonnegative edge length $\alpha : E\Gamma \to \mathbf{R}_+$, we define directed metric $D_{\Gamma,\alpha}$ on $V\Gamma$ as follows. For two nodes u, v, the distance $D_{\Gamma,\alpha}(u, v)$ is defined by the sum of edge-length $\alpha(e)$ over edges $e = pq$ such that the unique walk from u to v passes through pq in order $u \to p \to q \to v$. Namely $D_{\Gamma,\alpha}$ does not count the edge-length of edges with the opposite direction. A *subtree* of Γ is a subgraph whose underlying undirected graph is a tree. For a directed distance μ on S, an *oriented-tree realization* $(\Gamma, \alpha; \{F_s\}_{s\in S})$ is a triple of an oriented tree Γ, a nonnegative edge-length α, and a family $\{F_s\}_{s\in S}$ of subtrees indexed by S such that

$$\mu(s,t) = D_{\Gamma,\alpha}(F_s, F_t) \quad (s, t \in S).$$

Deletion of an edge $e = uv$ in Γ decomposes Γ into two connected components Γ'_e, Γ''_e so that Γ'_e contains u. This yields a partial cut (A_e, B_e) of S by $A_e := \{s \in S \mid F_s \text{ belongs to } \Gamma'_e\}$ and $B_e := \{s \in S \mid F_s \text{ belongs to } \Gamma''_e\}$. From definition of $D_{\Gamma,\alpha}$, one can easily see

$$\mu = \sum_{e \in E\Gamma} \alpha(e) \delta_{A_e,B_e}. \tag{3.2}$$

3.2 Proof of Theorem 3.1 (2)

Suppose $\dim \bar{Q}^{slim}_\mu \leq 1$. Then we can take a slimmed section R represented as a union of one-dimensional faces of $(Q^{slim}_\mu)^+$; see the proof of Lemma 2.5. By (3) and (4) in (2.7), each segment in R is isometric to a segment in $(\mathbf{R}, D^+_\infty)$. Since R is contractible (Corollary 2.6), the 1-skeleton graph Γ of R is a tree. Orient this 1-skeleton graph Γ so that for each edge pq (segment $[p, q]$), p is oriented to $q \Leftrightarrow D_\infty(p, q) > 0$ (and $D_\infty(q, p) = 0$). Also let $\alpha(pq) := D_\infty(p, q)$ for (oriented) edge $pq \in E\Gamma$. Then we obtain an oriented tree Γ with edge-length α. Let $\text{Vert}R$ be the set of vertices (endpoints of segments) of R. Since R is geodesic (Corollary 2.6 (1)), $(\text{Vert}R, D_\infty)$ is isometric to $(V\Gamma, D_{\Gamma,\alpha})$. For $s \in S$, let F_s be the subgraph induced by R_s (well-defined since R_s is a subcomplex of R). Since R_s is also contractible (Corollary 2.6 (1)), F_s is a subtree. Summarizing these facts together with Corollary 2.6 (2), we can conclude that $(\Gamma, \alpha; \{F_s\}_{s\in S})$ is an oriented-tree realization of μ.

I. We first prove the following min-max relation:

$$\begin{aligned} \text{MFP}^*(\mu; G, S, c) &= \text{FLP}^*(\text{Vert}R; G, S, c) \\ &= \min\left\{ \sum_{xy \in EG} c(xy) D_{\Gamma,\alpha}(\rho(x), \rho(y)) \;\middle|\; \rho : VG \to V\Gamma, \rho(s) \in VF_s \ (s \in S) \right\}. \end{aligned} \tag{3.3}$$

This means that FLP becomes a *discrete location problem* on Γ. By construction of Γ, it suffices to show the first equality, that is, there is an optimal map $\rho^* : VG \to R$ for FLP with $\rho^*(VG) \subseteq \mathrm{Vert}R$. Take any optimal map $\rho : VG \to R$. Suppose that there is an interior point p^* of some segment $[p,q]$ in R with $\rho^{-1}(p^*) \neq \emptyset$. Take a sufficiently small positive $\epsilon > 0$. Increase ϵ until $p^* + \epsilon(p-q) = p$ or $p^* - \epsilon(p-q) = q$ or $\rho^{-1}(p^* - \epsilon(p-q)) \neq \emptyset$ or $\rho^{-1}(p^* + \epsilon(p-q)) \neq \emptyset$. Let $\rho_+, \rho_- : VG \to R$ be defined by

$$\rho_\pm(x) = \begin{cases} p^* \pm \epsilon(p-q) & \text{if } \rho(x) = p^*, \\ \rho(x) & \text{otherwise,} \end{cases} \qquad (x \in VG).$$

Then both ρ_+ and ρ_- are feasible. Since $D_\infty(p,q) = D_\infty(p,r) + D_\infty(r,q)$ for $r \in [p,q]$ and R is geodesic, the following holds:

$$D_\infty(\rho(x),\rho(y)) = \frac{D_\infty(\rho_+(x),\rho_+(y)) + D_\infty(\rho_-(x),\rho_-(y))}{2} \quad (x,y \in VG).$$

Therefore both ρ_+ and ρ_- are optimal. For at least one of ρ_+, ρ_-, say ρ_+, the number of points $p \in R \setminus \mathrm{Vert}R$ with $(\rho_+)^{-1}(p) \neq \emptyset$ decreases. Let $\rho := \rho_+$. We can repeat this procedure until $\rho(VG) \subseteq \mathrm{Vert}R$. This proves claim (3.3).

II. Second we derive the following min-cut expression:

$$\mathrm{MFP}^*(\mu; G, S, c) = \sum_{e \in E\Gamma} \alpha(e) \min\{c(\partial X) \mid A_e \subseteq X \subseteq VG \setminus B_e\}. \tag{3.4}$$

($\leq$) follows from LHS $\leq \sum_{e\in E\Gamma} \alpha(e)\mathrm{MFP}^*(\delta_{A_e,B_e}; G,S,c) =$ RHS, where the inequality follows from (3.2), and the equality follows from the max-flow min-cut theorem (3.1). Let $\rho^* : VG \to V\Gamma$ be an optimal map in (3.3). Let d^* be the metric on VG defined by $d^*(x,y) = D_{\Gamma,\alpha}(\rho^*(x), \rho^*(y))$ for $x,y \in VG$. Then d^* has an oriented-tree realization $(\Gamma, \alpha; \{\rho(x)\}_{x\in VG})$. Again the deletion of edge e yields a cut (X_e, Y_e) of VG with $A_e \subseteq X_e \subseteq V \setminus B_e$, and $d^* = \sum_{e\in E\Gamma} \alpha(e)\delta_{X_e,Y_e}$. Thus $\mathrm{MFP}^*(\mu; G,S,c) = \sum_{xy\in EG} c(xy)d^*(x,y) = \sum_{e\in E\Gamma} \alpha(e) \sum_{xy \in EG} c(xy)\delta_{X_e,Y_e}(x,y) = \sum_{e\in E\Gamma}\alpha(e)c(\partial X_e)$.

III. Finally, we show the existence of an integral optimal multiflow. We use the splitting-off technique. By multiplying edges, we may assume that each edge has unit capacity. For a pair (xy, yz) of consecutive edges, the splitting-off operation is to delete xy and yz and add a new edge from x to z (of unit capacity). If the splitting-off operation does not decrease the optimal multiflow value, then from any optimal multiflow in the new network after the splitting-off we obtain an optimal multiflow in the initial network, and we can apply the inductive argument (on the number of edges). Consider any optimal (fractional) multiflow $f = (\mathcal{P}, \lambda)$. Suppose that there is a pair (xy, yz) of consecutive edges such that some path in $\mathcal{P}$ with nonzero flow-value passes through xy, yz in order. If such a pair does not exist, then f is already an integral multiflow. We show that the splitting-off at (xy, yz) is successful. Suppose that the splitting-off decreases the optimal flow-value. By (3.4), there are $e \in E\Gamma$ and X^* attaining the minimum of $\min\{c(\partial X) \mid A_e \subseteq X \subseteq VG \setminus B_e\}$ such that $(*)$ $x, z \in X^* \not\ni y$ or $y \in X^* \not\ni x, z$. Since f is an optimal multiflow for weight δ_{A_e,B_e}, i.e., a maximum (single commodity) (A_e, B_e)-flow, each path in $\mathcal{P}$ (with nonzero flow-value) must meet ∂X^* at most once. This contradicts $(*)$.

3.3 Combinatorial min-max relations

We have already shown that if $\dim \bar{Q}_\mu^{slim} \leq 1$, then we obtain an oriented-tree realization of μ by $\bar{Q}_\mu^{slim}$, and the min-max relation (3.3) from this realization. The next theorem states that if μ is realized by an oriented tree, then one can get such a min-max relation directly (without calculating $\bar{Q}_\mu^{slim}$). Let $\mathrm{IMFP}^*(\mu; G, S, c)$ denote the maximum flow-value with respect to μ over all integral multiflows in (G, S, c).

Theorem 3.2. *Let μ be a directed distance on S having an oriented-tree realization $(\Gamma, \alpha; \{F_s\}_{s\in S})$, and let (G, S, c) be an inner Eulerian network such that the Eulerian condition is fulfilled by each terminal s with F_s being neither a single node nor a directed path. Then the following relation holds:*

$$
\begin{aligned}
(3.5)\qquad \mathrm{MFP}^*(\mu; G, S, c) &= \mathrm{IMFP}^*(\mu; G, S, c)\\
&= \min\left\{ \sum_{xy\in EG} c(xy) D_{\Gamma,\alpha}(\rho(x), \rho(y)) \;\middle|\; \rho : VG \to V\Gamma,\ \rho(s) \in VF_s\ (s \in S) \right\}\\
&= \sum_{e\in E\Gamma} \alpha(e) \min\{c(\partial X) \mid A_e \subseteq X \subseteq VG \setminus B_e\},
\end{aligned}
$$

where (A_e, B_e) is a partial cut on S determined by the deletion of edge $e \in E\Gamma$.

The proof uses the next proposition, which says that $(Q_\mu^{slim})^+$ is (essentially) a geometric realization of an oriented-tree realization $(\Gamma, \alpha; \{F_s\}_{s\in S})$.

Proposition 3.3. *Suppose that μ has an oriented-tree realization $(\Gamma, \alpha; \{F_s\}_{s\in S})$ so that $\{F_s\}_{s\in S}$ contains all single-node subtrees. Let $S_0 \subseteq S$ consist of elements s such that F_s is a single node v_s. Then the following holds:*

1. $(Q_\mu^{slim})^+ = \bigcup\{[\mu_s, \mu_t] \mid s, t \in S_0, v_s v_t \in E\Gamma\}$.
2. $(Q_\mu^{slim})_s^+ = \bigcup\{[\mu_t, \mu_u] \mid t, u \in S_0, v_t v_u \in EF_s\}$ *for $s \in S$.*
3. $(Q_\mu^{slim})^+$ *itself is a slimmed section.*
4. $(Q_\mu)_s$ *has no fat if F_s is a single node or a directed path.*

See Section 2.1 B for definition of μ_s. The proof is a routine verification, but rather technical. So the proof is given in the end of this subsection. In (4) the converse (only-if part) also holds. However we omit the proof, which is also a lengthy verification.

Assuming Proposition 3.3, we complete the proof of Theorem 3.2. Suppose that μ is realized by $(\Gamma, \alpha; \{F_s\}_{s\in S})$. We can add isolated terminals to (G, S, c) so that $\{F_s\}_{s\in S}$ includes all single-node subtrees. Thus we may assume that $(\Gamma, \alpha; \{F_s\}_{s\in S})$ fulfills the hypothesis in Proposition 3.3. Consider a slimmed section $R = (Q_\mu^{slim})^+$. Then the 1-skeleton graph of R coincides with Γ. Hence we can apply the arguments (e.g., (3.3), (3.4)) in the previous subsection.

We give characterizations of a class of distances μ with $\dim \bar{Q}_\mu^{slim} \leq 1$. Two partial cuts (A, B) and (A', B') are said to be *laminar* if $A \subseteq A', B \supseteq B'$ or $A \subseteq B', B \supseteq A'$ or $A \supseteq A', B \subseteq B'$ or $A \supseteq B', B \subseteq A'$. A family $\mathcal{A}$ of partial cuts is said to be *laminar* if every pair in $\mathcal{A}$ is laminar.

Theorem 3.4. *For a directed distance μ on S, the following conditions are equivalent:*

1. $\dim \bar{Q}_\mu^{slim} \leq 1$.
2. *μ has an oriented-tree realization.*
3. *There are a laminar family $\mathcal{A}$ of partial cuts on S and a positive weight $\alpha : \mathcal{A} \to \mathbf{R}_+$ such that*
$$\mu = \sum_{(A,B)\in\mathcal{A}} \alpha(A, B)\delta_{A,B}.$$

Proof. We have already seen (1) $\Rightarrow$ (2) in Section 3.2. Theorem 4.1 (2) in Section 4 says that if $\dim \bar{Q}_{\mu}^{slim} \geq 2$ then there is no integer k such that μ-MFP has a $1/k$-integral multiflow for every Eulerian network. Therefore, by Theorem 3.2, if μ has an oriented-tree realization, then $\dim \bar{Q}_{\mu}^{slim} \leq 1$ necessarily holds. Thus we have (2) $\Rightarrow$ (1). The equivalence (2) $\Leftrightarrow$ (3) is not difficult, and is essentially obtained by [4] (in an undirected version). $\square$

Directed multiflow locking theorem. Let $\mathcal{A}$ be a set of partial cuts on terminal set S in a network. We say that a multiflow f *locks* $\mathcal{A}$ if f is simultaneously a maximum (A, B)-flow for all partial cuts (A, B) in $\mathcal{A}$. In the case where $\mathcal{A}$ is laminar, there are an oriented-tree $\mathit{\Gamma}$ and a family $\{F_s\}_{s \in S}$ of subtrees such that $\mathcal{A}$ coincides with the set $\{(A_e, B_e)\}_{e \in E\mathit{\Gamma}}$ of partial cuts on S. Consider distance $\mu := \sum_{(A,B) \in \mathcal{A}} \delta_{A,B}$. Then μ is realized by $(\mathit{\Gamma}, 1; \{F_s\}_{s \in S})$. Here F_s is a directed path if (and only if) there is no pair $(A, B), (A', B') \in \mathcal{A}$ with $s \notin A \cup B \cup A' \cup B'$ and $A \subseteq B', A' \subseteq B$ or $B \subseteq A', B' \subseteq A$. Apply Theorem 3.2 to μ. From the last equality in (3.5), an optimal multiflow is necessarily optimal to $\delta_{A,B}$-MFP for each $(A, B) \in \mathcal{A}$; see the argument after (3.4). This implies the following:

Theorem 3.5. *Let $\mathcal{A}$ be a laminar family of partial cuts on S, and let (G, S, c) be an inner Eulerian network. If the Eulerian condition is fulfilled by each terminal s having a pair $(A, B), (A', B') \in \mathcal{A}$ with $s \notin A \cup B \cup A' \cup B'$ and $A \subseteq B', A' \subseteq B$ or $B \subseteq A', B' \subseteq A$, then there is an integral multiflow locking $\mathcal{A}$.*

This includes Ibaraki-Karzanov-Nagamochi's result for laminar cuts.

Theorem 3.6 ([9, Theorem 5])**.** *Let $\mathcal{A}$ be a laminar family of cuts on S. For every inner Eulerian network (G, S, c), there is an integral multiflow locking $\mathcal{A}$.*

0-1 distances and commodity graphs. Suppose the case where μ is $\{0, 1\}$-valued. In this case, μ can be identified with a *commodity graph* H by $st \in EH \Leftrightarrow \mu(s, t) = 1$. For a commodity graph H on S, let μ_H denote the corresponding 0-1 distance on S defined by $\mu_H(s, t) = 1 \Leftrightarrow st \in EH$. In the case where H is a complete digraph, Lomonosov and Frank independently established the following min-max relation:

Theorem 3.7 ([14, 3])**.** *Let H be a complete digraph on S. For every inner Eulerian network (G, S, c), we have*

$$\mathrm{MFP}^*(\mu_H; G, S, c) = \mathrm{IMFP}^*(\mu_H; G, S, c) = \sum_{s \in S} \min\{c(\partial X) \mid s \in X \subseteq VG \setminus (S \setminus s)\}.$$

This theorem can be regarded as a special case of Theorem 3.2. Indeed, the all-one distance is realized by a star with the sink (or source) as its center. So we can extend this theorem to a class of commodity graphs having oriented-tree realizations.

A *quasi-complete digraph* H is a simple digraph having a node subset T such that

(0) all edges are incident to T,

(1) the subgraph induced by T is a complete digraph, and

(2) all edges between T and $VH \setminus T$ leave T or enter T.

The node set T is said to be the *complete part*, and H is said to be *source-type* if the edges between T and $VH \setminus T$ enter T and is said to be *sink-type* otherwise. For a quasi-complete digraph H with complete part $T = \{x_1, x_2, \ldots, x_m\}$, the corresponding $\{0, 1\}$-valued distance μ_H has an oriented-tree realization by a star $\mathit{\Gamma}$ of m leaves $v_1, v_2, \ldots, v_m$

such that the center v_0 is a source if H is source-type, and is a sink if H is sink-type. Indeed, for $i = 1, 2, \ldots, m$, let R_{x_i} be the subtree consisting of one node v_i. For a node $s \in VH \backslash T$, if s is joined to $x_{j_1}, x_{j_2}, \ldots, x_{j_k}$, then let R_s be the subtree consisting of nodes $\{v_0, v_1, v_2, \ldots, v_m\} \setminus \{v_{j_1}, v_{j_2}, \ldots, v_{j_k}\}$. Then one can verify that $(\mathit{\Gamma}, 1; \{R_s\}_{s \in VH})$ is a required realization. In particular, each node s having at least $m-1$ edges is associated with a single node or a directed path in $\mathit{\Gamma}$. By Theorem 3.2, we have the following:

Theorem 3.8. *Let H be a quasi-complete digraph on S with complete part T, and let (G, S, c) be an inner Eulerian network such that the Eulerian condition is fulfilled by each terminal s incident to at most $|T| - 2$ edge in H. Then the following holds:*

$$\text{(3.6)} \qquad \mathrm{MFP}^*(\mu_H; G, S, c) = \mathrm{IMFP}^*(\mu_H; G, S, c)$$
$$= \begin{cases} \displaystyle\sum_{s \in T} \min\{c(\partial X) \mid s \in X \subseteq VG \setminus N_H(s)\} & \text{if } H \textit{ is sink-type,} \\ \displaystyle\sum_{s \in T} \min\{c(\partial X) \mid N_H(s) \subseteq X \subseteq VG \setminus s\} & \text{if } H \textit{ is source-type,} \end{cases}$$

where $N_H(s)$ is the set of nodes incident to s in H.

A *multipartite extension* of a graph H is a graph obtained by replacing each node v by a node subset U_v and joining each pair $(x, y) \in U_v \times U_u$ exactly when $vu \in EH$. Trivially we can further extend this relation (3.6) to the case where H is a multipartite extension of a quasi-complete digraph (by super sink/source argument).

Also we easily see from Theorem 3.4 the following:

Proposition 3.9. *For a simple digraph H on S, the following conditions are equivalent:*

(a) $\dim \bar{Q}^{slim}_{\mu_H} \leq 1$.

(b) *H is a multipartite extension of a quasi-complete digraph.*

Proof of Proposition 3.3. Take an arbitrary $s \in S_0$. We first claim $\mu_s \in (Q^{slim}_\mu)^+$. Since F_s is a single node v_s, we have $\mu_s(t^c) + \mu_s(u^r) = \mu(t, s) + \mu(s, u) = D_{\mathit{\Gamma},\alpha}(F_t, v_s) + D_{\mathit{\Gamma},\alpha}(v_s, F_u) \geq D_{\mathit{\Gamma},\alpha}(F_t, F_u) = \mu(t, u)$ for $t, u \in S$. Thus $\mu_s \in P^+_\mu$. Next we give a description of $K(\mu_s)$. Delete v_s from $\mathit{\Gamma}$. Let $\mathit{\Gamma}_1, \mathit{\Gamma}_2, \ldots, \mathit{\Gamma}_k$ be the resulting connected components. For $i = 1, 2, \ldots, k$, let U_i be the set of elements $t \in S$ such that F_t belongs to $\mathit{\Gamma}_i$. Let W be the set of elements $t \in S$ such that F_t contains v_s. Then $\{W, U_1, U_2, \ldots, U_k\}$ is a partition of S. A pair $(t^c, u^r) \in S^c \times S^r$ has an edge in $K(\mu_s)$ if and only if a shortest path from F_t to F_u can pass through the node v_s. We remark that tracing an edge in reverse direction takes zero length. Then we see the following:

(a) Pair $(u^c, t^r) \in U_i{}^c \times U_j{}^r$ has an edge if and only if $i \neq j$.

(b) Each pair $(u^c, t^r) \in W^c \times W^r$ has an edge.

(c) s^c is incident to each element in S^r and s^r is incident to each element in S^c.

So there is no isolated node, and thus we have $\mu_s \in Q^+_\mu$. By (c), μ_s is not a fat. Thus $\mu_s \in (Q^{slim}_\mu)^+$ and in particular $\mu_s \in Q^+_{\mu,W}$ by (a,b). Also we see:

(d) For $1 \leq i \leq k$ there is $t \in U_i \cap S_0$ such that either $\mu_s(t^c) = 0$ or $\mu_s(t^r) = 0$.

Indeed, by assumption, there is $t \in S_0$ such that v_t is a node in Γ_i incident to v_s. Then $t \in U_i$, and $\mu_s(t^c) = 0$ if $v_s v_t \in E\Gamma$ and $\mu_s(t^r) = 0$ if $v_t v_s \in E\Gamma$. Next we claim:

(3.7) If a face F of $(Q_\mu^{slim})^+$ contains μ_s, then $F = [\mu_s, \mu_t]$ for some $t \in S_0$ with $v_s v_t \in E\Gamma$ or $v_t v_s \in E\Gamma$.

If true, then we obtain the first statement (1) (since $(Q_\mu^{slim})^+$ is connected). Perturb μ_s into p so that $p \in (Q_\mu^{slim})^+$ and $EK(p) \subseteq EK(\mu_s)$ (i.e., p belongs to a face containing μ_s). Let X^- be the set of nodes $u \in S^{cr}$ with $p(u) < \mu_s(u)$, and let X^+ be the set of nodes $u \in S^{cr}$ with $p(u) > \mu_s(u)$. Recall (2.8). Necessarily $X^+ = N_{\mu_s}(X^-)$; otherwise there is an isolated node in $K(p)$. We claim

($*$) $X^- = \bigcup_{j \in I} U_j{}^c$ or $X^- = \bigcup_{j \in I} U_j{}^r$ for some $I \subseteq \{1, 2, \ldots, k\}$.

Suppose that both $U_j{}^c \cap X^-$ and $U_j{}^c \setminus X^-$ are nonempty. Since $N_{\mu_s}(U_j{}^c \cap X^-) \setminus W^r = N_{\mu_s}(U_j{}^c \setminus X^-) \setminus W^r$ by (a), $W^r \subseteq N_{\mu_s}(U_j{}^c \cap X^-)$ implies that $U_j{}^c \setminus X^-$ is isolated. So $W^r \setminus N_{\mu_s}(U_j{}^c \cap X^-)$ is nonempty, and has an edge incident to $U_j{}^c \setminus X^-$. Let $W_0 \subseteq W$ with $W_0{}^r := W^r \setminus N_{\mu_s}(U_j{}^c \cap X^-)$. Necessarily each F_t for $t \in W_0$ includes edge e joining v_s and Γ_j (otherwise t^r is incident to all elements in $U_j{}^c$), and moreover e leaves v_s (otherwise there is no edge between $W_0{}^r$ and $U_j{}^c$). By this property, there is no edge joining $W_0{}^c$ and $U_j{}^r$ in $K(\mu_s)$ (and in $K(p)$). Thus p is a proper fat relative to W_0; a contradiction. Also $U_i{}^c \cup U_j{}^r \subseteq X^-$ is impossible by (a,d).

We may suppose $X^- = \bigcup_{j \in I} U_j{}^c$. We show $I = \{i\}$ for some i. Suppose true. Then we can see $p = \mu_s + \epsilon(-\mathbf{1}_{U_i{}^c}, \mathbf{1}_{N_{\mu_s}(U_i{}^c)})$ for some $\epsilon > 0$. By (d), there is $t \in U_i \cap S_0$ with $v_t v_s \in E\Gamma$ and $p \in [\mu_s, \mu_t]$, as required. Suppose $|I| \geq 2$. Then $S^r \setminus W^r \subseteq N_{\mu_s}(X^-) = X^+$ by (a), and $K(p)$ has no edge between $S^c \setminus X^-$ and X^+. This means that p is a proper fat relative to some $W' \subseteq W$; a contradiction.

In the argument above, we can see that the perturbed p never belongs to any degenerate set; so Q_μ has no degenerate set. This implies (3). The claim (2) can be verified in a straightforward manner.

(4). Let t be a terminal such that F_t is a single node or a directed path. Take any $s \in S_0$ with v_s belonging to F_t. Then μ_s belongs to $(Q_\mu^{slim})_t^+$. Again perturb μ_s into $p \in (Q_\mu)_t^+$. It suffices to show $p \in (Q_\mu^{slim})_t^+$. In the partition $\{W, U_1, U_2, \ldots, U_k\}$ for $K(\mu_s)$, t belongs to W. As above, consider X^-, X^+. Then $X^- = \bigcup_{j \in I} U_j{}^c$ or $X^- = \bigcup_{j \in I} U_j{}^r$ for some $I \subseteq \{1, 2, \ldots, k\}$. From the assumption that F_t is a single node or a directed path, one can see that t^r is incident to all nodes in S^c except $U_i{}^c$ for which Γ_i includes the tail of F_t, and that t^c is incident to all nodes in S^r except $U_j{}^r$ for which Γ_j includes the head of F_t. From this fact, either $I = \{i\}$ or $\{j\}$; otherwise edge $t^c t^r$ vanishes in $K(p)$ and this implies $p \notin (Q_\mu)_t^+$. Thus we can verify $p \in (Q_\mu^{slim})_t^+$ as above.

3.4 Case $\dim T_\mu \leq 1$: reduction to minimum cost circulation

Suppose $\dim T_\mu \leq 1$. In this case, T_μ is also a tree. Thus the argument in Section 3.2 is applicable. However, by (2.7) (5) T_μ is a path isometric to a segment in $(\mathbf{R}, D_\infty^+)$. Therefore by (2.5) (2) there is a family $\{[a_s, b_s] \mid s \in S\}$ of segments in $\mathbf{R}$ such that

$$\mu(s,t) = (a_t - b_s)_+ \quad (s, t \in S).$$

By using this expression, we show that μ-MFP is reducible to the minimum cost circulation. Let (G, S, c) be a network. For each terminal pair (s,t) with $\mu(s,t) = (a_t - b_s)_+ > 0$, add new edge (terminal edge) ts with edge-cost $-\mu(s,t)$. Then consider the minimum cost circulation problem on the new network; this is a relaxation of μ-MFP. As is well-known, there is an integral minimum cost circulation. This circulation can be decomposed into the sum of the incidence vectors for some (possibly repeating) cycles. If each cycle contains at most one terminal edge, then we obtain an integral optimal multiflow by deleting the terminal edge from each cycle. So suppose that there is a cycle C containing at least two terminal edges. Then C is the union of terminal edges $t_0t_1, t_2t_3, \ldots, t_{k-1}t_k$ and S-paths $P_{1,2}, P_{3,4}, \ldots, P_{k-2,k-1}, P_{k,0}$, where k is an odd integer, and $P_{i,i+1}$ is an (t_i, t_{i+1})-path. We claim

$$\sum_{i=0,2,4,\ldots,k-1} \mu(t_{i+1}, t_i) \leq \sum_{i=0,2,4,\ldots,k-1} \mu(t_{i-1}, t_i), \tag{3.8}$$

where we let $t_{-1} = t_k$. The LHS ($=: \mu(C)$) is the negative of the cost of the cycle C and the RHS is the total flow-value of S-paths $\{P_{i,i+1}\}_{i=1,3,5,\ldots}$ (with unit flow-values). Suppose that the claim (3.8) is true. By decomposing each cycle into S-paths as above, we obtain an integral multiflow f whose total flow-value $\mathrm{val}(\mu, f)$ is at least the negative of the total cost of the mincost relaxation problem. So f is optimal.

The claim (3.8) can be seen as follows. Move point x in $\mathbf{R}$ as $a_{t_0} \to b_{t_1} \to a_{t_2} \to b_{t_3} \to \cdots \to b_{t_k} \to a_{t_0}$. In each odd step, the point x moves in the negative direction since $a_{t_i} > b_{t_{i+1}}$. In particular the total move over odd steps coincides with $\mu(C)$. Since the point x returns to the initial point, $\mu(C)$ is at most the total move in the positive direction over even steps, which equals the RHS in (3.8).

4 Unbounded fractionality

The integrality theorem (Theorem 3.1) in the previous section is best possible. The goal of this section is to establish the unbounded fractionality property:

Theorem 4.1. *Let μ be a directed distance on S.*

(1) *If $\dim T_\mu \geq 2$, then there is no positive integer k such that μ-MFP has a $1/k$-integral optimal multiflow for every network (G, S, c).*

(2) *If $\dim \bar{Q}_\mu^{slim} \geq 2$, then there is no positive integer k such that μ-MFP has a $1/k$-integral optimal multiflow for every totally Eulerian network (G, S, c).*

In the following, the edge set of a complete digraph (without loops) on a set V is denoted by E_V. We regard a function $g : V \times V \to \mathbf{R}_+$ with zero diagonals $g(x, x) = 0$ for $x \in V$ as $E_V \to \mathbf{R}_+$; we simply denote $g(x, y)$ by $g(xy)$.

We utilize Edmonds-Giles' lemma for rational polyhedra; see [15, Section 22.1]:

(4.1) For an integer $k > 0$, a rational polyhedron $P \subseteq \mathbf{R}^n$ is $1/k$-integral if and only if $\min\{\langle c, x\rangle \mid x \in P\}$ is a $1/k$-integer for each integral vector $c \in \mathbf{Z}^n$ for which the minimum is finite.

Here a polyhedron P is said be $1/k$-*integral* if each face of P contains a $1/k$-integral vector, and $\langle \cdot, \cdot \rangle$ denotes the standard inner product in $\mathbf{R}^n$. For a finite set $V \supseteq S$, consider the following two unbounded polyhedra:

$$\begin{aligned} \mathcal{D}_{\mu,V} &:= \{d\text{: metric on } V \mid d(st) \geq \mu(st)\ (st \in E_S)\} + \mathbf{R}_+^{E_V}, \\ \mathcal{D}'_{\mu,V} &:= \mathcal{D}_{\mu,V} + \mathcal{L}, \\ &\quad\ \text{where } \mathcal{L} := \{l \in \mathbf{R}^{E_V} \mid l(C) = 0 \text{ (all cycles } C \text{ in } V)\}. \end{aligned}$$

Note that $\mathcal{D}_{\mu,V}$ is pointed, and $\mathcal{D}'_{\mu,V}$ is not pointed. Then $\min\{\langle c,d\rangle \mid d \in \mathcal{D}_{\mu,V}\}$ is finite if and only if c is nonnegative, and if finite, then it equals $\mathrm{MFP}^*(\mu;(V,E_V),S,c)$. Also $\min\{\langle c,d\rangle \mid d \in \mathcal{D}'_{\mu,V}\}$ is finite if and only if $((V,E_V),S,c)$ is totally Eulerian, and if finite, then it equals $\mathrm{MFP}^*(\mu;(V,E_V),S,c)$. Hence it suffices to show:

Proposition 4.2. *Let μ be a directed distance on S, and let k be any positive integer.*

(1) *If $\dim T_\mu \geq 2$, then $\mathcal{D}_{\mu,V}$ is not $1/k$-integral for some $V \supseteq S$.*

(2) *If $\dim \bar{Q}^{slim}_\mu \geq 2$, then $\mathcal{D}'_{\mu,V}$ is not $1/k$-integral for some $V \supseteq S$.*

Indeed, if a $1/k$-integral optimal multiflow always exists, then the optimal value is always $1/k$-integral, and $\mathcal{D}_{\mu,V}$ is $1/k$-integral for all V by (4.1). The rest of this section is devoted to the proof of this proposition. We note the following relation for two metrics d, d' on V, which follows from the cycle decomposition of a circulation.

(4.2) $d \equiv d' \mod \mathcal{L}$ if and only if $d(xy) = d'(xy) - p(x) + p(y)$ $(xy \in E_V)$ for some $p : V \to \mathbf{R}$.

4.1 Preliminary: minimal and extreme metrics

We begin with preliminary arguments. A metric $d \in \mathcal{D}_{\mu,V}$ is said to be *minimal* if there is no other metric $d' \in \mathcal{D}_{\mu,V}$ with $d' \neq d$ and $d' \leq d$, and is said to be $\mathcal{C}$*-minimal* if there is no other metric $d' \in \mathcal{D}_{\mu,V}$ with $d' \not\equiv d \mod \mathcal{L}$ and $d'(C) \leq d(C)$ for all cycles C.

We first give characterizations of minimal and $\mathcal{C}$-minimal metrics. Let d be a metric on S. Then d defines the equivalence relation on S by $x \sim_d y \overset{\text{def}}{\iff} d(xy) = d(yx) = 0$. Let $[x]$ denote the equivalence class including x, and let $[xy]$ denote the set of edges from $[x]$ to $[y]$. An edge $xy \in E_S$ is said to be *extremal* if there is no edge $st \in E_S \setminus [xy]$ with $d(st) = d(sx) + d(xy) + d(yt)$. An edge xy is extremal if and only if $x'y' \in [xy]$ is extremal. So a class $[xy]$ is said to be *extremal* if xy is extremal. Let $H_{\mu,d}$ be the directed graph on S with edge set $EH_{\mu,d} = \{st \in E_S \mid d(st) = \mu(st)\}$.

Lemma 4.3. *Let d be a directed metric in $\mathcal{D}_{\mu,S}$.*

(1) *d is minimal if and only if every extremal class meets an edge in $H_{\mu,d}$.*

(2) *d is $\mathcal{C}$-minimal if and only if every extremal class meets a cycle in $H_{\mu,d}$.*

We note that $d(xy) = d(x'y')$ for $x'y' \in [xy]$ and $xy, yx \in H_{\mu,d}$ if $[x] = [y]$. In particular xy is never extremal if $[x] = [y]$ (and $d \neq 0$). In most cases, we may consider the case where each class consists of one edge.

Proof. (1). If part: By condition we cannot decrease d on extremal classes. Consequently we cannot decrease d on non-extremal classes by the triangle inequality. Only-if part: suppose that some extremal class $[xy]$ fulfills $d(uv) > \mu(uv)$ for $uv \in [xy]$. We can decrease d on $[xy]$ with keeping triangle inequality.

(2). For $p : S \to \mathbf{R}$, let $d * p$ be defined by $(d * p)(xy) = d(xy) - p(x) + p(y)$ for $xy \in E_S$. By definition and (4.2), we see:

($*$1) d is $\mathcal{C}$-minimal if and only if $d * p$ is minimal for every $p : S \to \mathbf{R}$ with $d * p \in \mathcal{D}_{\mu,S}$.

($*$2) The set of extremal edges of d is the same as that of $d * p$.



Figure 6: Triangulations

First we show the if part (of (2)). Take any $p : S \to \mathbf{R}$ with $d * p \in \mathcal{D}_{\mu,S}$, and take any extremal class $[st]$ of $d * p$. Since $[st]$ is also an extremal class for d by $(*2)$, $[st]$ meets a cycle C in $H_{\mu,d}$. Therefore $\sum_{xy \in C}(d * p)(xy) = \sum_{xy \in C} d(xy) = \sum_{xy \in C} \mu(xy)$. By $d(xy) - p(x) + p(y) = (d * p)(xy) \geq \mu(xy)$, p is constant on C; in particular $(d * p)(xy) = d(xy) = \mu(xy)$ on C. This means that $d * p$ is minimal by (1). Thus d is $\mathcal{C}$-minimal by $(*1)$.

We show the only-if part. Suppose that some extremal class $[st]$ does not meet any cycle. In this case, there is a node subset $U \subseteq S$ such that $[s] \subseteq U$, $[t] \subseteq S \setminus U$, and there is no edge entering U (consider the strong component decomposition of $H_{\mu,d}$). For a sufficiently small $\epsilon > 0$, let $p : S \to \mathbf{R}$ be defined by $p(u) = 0$ for $u \in U$ and $p(u) = \epsilon$ for $u \notin U$. Then $d * p \in \mathcal{D}_{\mu,S}$ and $(d * p)(uv) = d(uv) + \epsilon > \mu(uv)$ for $uv \in [st]$. Thus $d * p$ is not minimal, and d is not $\mathcal{C}$-minimal. $\square$

Second we recall the notion of extreme metrics. A metric d on a finite set V is called *extreme* if d lies on an extreme ray the polyhedral cone $\mathcal{M}_V$ formed by all metrics on V. Also d is $\mathcal{C}$-*extreme* if the projection $d/\mathcal{L}$ is extreme in $\mathcal{M}_V/\mathcal{L}$. We give a family of extreme metrics. Consider the refinement sequence of triangulations of a plane triangle $\{(x, y) \in \mathbf{R}^2 \mid 0 \leq y \leq x \leq 1\}$ in $(\mathbf{R}^2, D_\infty^+)$ by congruent triangles, as in Figure 6. Let γ_n be the metric obtained by restricting $(\mathbf{R}^2, D_\infty^+)$ to the vertex set of the n-th triangulation.

Lemma 4.4. *γ_n is extreme and $\mathcal{C}$-extreme.*

Proof. One can easily verify that γ_1 is extreme. Suppose $\gamma_n = d' + d''$ for some metrics d', d''. We show $\gamma' = \alpha\gamma_n$ for some positive α. We observe that the restriction of γ_n to each triangle is isometric to $(1/n)\gamma_1$, which is extreme. Since the triangulation is *connected*, we can take a common positive α such that $d'(pq) = \alpha\gamma_n(pq)$ for p, q in any triangle. For an arbitrary pair p, q of vertices, there is a path $p = p_1, p_2, \ldots, p_m = q$ lying on the triangulation graph such that $\gamma_n(pq) = \sum_{i=1}^{m-1} \gamma_n(p_i p_{i+1})$. By $\gamma_n = d' + d''$, this equality must hold for d'. Hence, we have $d'(pq) = \sum_i d'(p_i p_{i+1}) = \alpha \sum_i \gamma_n(p_i p_{i+1}) = \alpha\gamma_n(pq)$. Thus we have $d' = \alpha\gamma_n$.

Next we consider the $\mathcal{C}$-extremality. As above, the $\mathcal{C}$-extremality of γ_n $(n \geq 2)$ reduces to that of γ_1 by the following observations: For an arbitrary cycle C there is a cycle C' in the graph with $d(C) = d(C')$, and for an arbitrary cycle C' in the graph there are cycles $C_1, C_2, \ldots, C_m$ each of which belongs to a triangle such that $d(C') = \sum_{i=1}^{m} \pm d(C_i)$. The $\mathcal{C}$-extremality of γ_1 also follows from a routine calculation, and is left to readers. *Sketch:* suppose $\gamma_1 \equiv d' + d'' \mod \mathcal{L}$. By (4.2), $\gamma_1(xy) + \gamma_1(yz) = \gamma_1(xz)$ implies the same equality for $d' + d''$, which in turn implies the same equality for d'. By using it, we can show $d'(C) = \alpha\gamma_1(C)$ for $\alpha := d'(xy) + d'(yx)$. $\square$

4.2 Proof (metric case)

Suppose that μ is a metric. In this case, $\mathcal{D}_{\mu,V}$ is represented as

$$\mathcal{D}_{\mu,V} = \{d\text{: metric on } V \mid d(st) = \mu(st) \ (st \in E_S)\} + \mathbf{R}_+^{E_V}.$$

Recall the notions in Section 2.1 C. Then, metric d is minimal in $\mathcal{D}_{\mu,V}$ if and only if d is a tight extension of μ. Also d is $\mathcal{C}$-minimal in $\mathcal{D}_{\mu,V}$ if and only if d is a cyclically tight extension of μ.

We first prove Proposition 4.2 (2). $\bar{Q}_\mu^{slim} = \bar{Q}_\mu$ by Proposition 2.3. We can take a balanced section R in Q_μ^+ containing a 2-dimensional face F, which is isometric to a polygon in $(\mathbf{R}^2, D_\infty^+)$ by (2.7) (3). Therefore we can take a subset U in F whose metric induced by D_∞^+ is isometric to $\beta\gamma_n$ for some $\beta > 0$. Fix an integer $k > 0$. By (2.6) (2), for an arbitrary integer $n > 0$, we can take a cyclically tight extension d on V having $\beta\gamma_n$ as a submetric. Take a sufficiently large n. Since $d/\mathcal{L}$ belongs to a bounded face of $\mathcal{D}_{\mu,V}/\mathcal{L}$, we can decompose d into a convex combination of cyclically tight extensions $d_1, d_2, \ldots, d_m$ in modulo $\mathcal{L}$ such that $d_i/\mathcal{L}$ is an extreme point in $\mathcal{D}_{\mu,V}/\mathcal{L}$ for $i = 1, 2, \ldots, m$. Since d has $\beta\gamma_n$ as a submetric and γ_n is $\mathcal{C}$-extreme, some d_i has a submetric γ with $\gamma \equiv \alpha\beta\gamma_n \mod \mathcal{L}$ for some $\alpha > 0$. Therefore $d_i(pq) + d_i(qp) = \alpha\beta/n$ for some $pq \in E_V$. Since d_i is cyclically tight, it is embedded into (Q_μ^+, D_∞). Therefore $\alpha\beta$ is bounded by the diameter of Q_μ^+ (bounded set). Since n is sufficiently large, we have $\alpha\beta/n < 1/k$. Hence face $d_i + \mathcal{L}$ has no $1/k$-integer vector.

Proposition 4.2 (1) can be shown in a similar manner. Since T_μ has a 2-dimensional face, we can take a tight extension d having $\beta\gamma_n$ as a submetric. Since $\mathcal{D}_{\mu,V}$ is pointed and d belongs to a bounded face in $\mathcal{D}_{\mu,V}$ by minimality, we can decompose d into a convex combination of extreme points in $\mathcal{D}_{\mu,V}$. For a sufficiently large n, one of the summands is not $1/k$-integral as above.

4.3 Proof (general case)

Suppose that μ is not a metric. For a distance g on S and a subset $U \subseteq S$, the restriction of g to U is denoted by g_U.

Lemma 4.5. *Let d be a directed metric in $\mathcal{D}_{\mu,V}$.*

(1) *If d_S is minimal in $\mathcal{D}_{\mu,S}$ and d is a tight extension of d_S, then d is minimal in $\mathcal{D}_{\mu,V}$.*

(2) *If d_S is $\mathcal{C}$-minimal in $\mathcal{D}_{\mu,S}$ and d is a cyclically tight extension of d_S, then d is $\mathcal{C}$-minimal in $\mathcal{D}_{\mu,V}$.*

Proof. (1) is obvious from definition. (2) is not so obvious. We utilize Lemma 4.3 by extending μ to $\bar{\mu} : E_V \to \mathbf{R}_+$ by $\bar{\mu}_S := \mu$ and $\bar{\mu}(xy) := 0$ for $xy \notin E_S$. Take an extremal class $[xy]$ of d. We show $[xy] \cap E_S \neq \emptyset$. If true, then $[xy] \cap E_S$ is also an extremal class in d_S and meets a cycle of H_{μ,d_S} by the $\mathcal{C}$-minimality of d_S (Lemma 4.3 (2)); this cycle also belongs to $H_{\bar{\mu},d}$.

Extend d_S to $\bar{d} : E_V \to \mathbf{R}_+$ by $\bar{d}_S := d_S$ and $\bar{d}(xy) := 0$ for $xy \notin E_S$. Since d is a cyclically tight extension of d_S, d is $\mathcal{C}$-minimal in $\mathcal{D}_{\bar{d},V}$. By Lemma 4.3 (2), $[xy]$ meets a cycle C in $H_{\bar{d},d}$. If this cycle belongs to $E_V \setminus E_S$, then by $d(uv) = \bar{d}(uv) = 0$ for $uv \in C$ the triangle equality we have $[x] = [y]$ and thus $[xy]$ is never extremal; a contradiction. Therefore C meets distinct nodes in S. By the triangle inequality and $\bar{d} = 0$ on $E_V \setminus E_S$ we may assume that C includes a path (u, x, y, v) with distinct $u, v \in S$. In particular $d(uv) = d(ux) = d(xy) = d(yv) = 0$. Since $d(uv) = d(ux) + d(xy) + d(yv)$ and xy is extremal, $d(xu) = d(vy) = 0$, and thus $([x], [y]) = ([u], [v])$. So $[xy] \cap E_S \neq \emptyset$. □

Our final goal is the following:

Lemma 4.6. *Let μ be a directed distance on S.*

1. *If $\dim T_\mu \geq k$, then there is a minimal metric d in $\mathcal{D}_{\mu,S}$ with $\dim T_d \geq k$.*
2. *If $\dim \bar{Q}^{slim}_\mu \geq 2$, then there is a $\mathcal{C}$-minimal metric d in $\mathcal{D}_{\mu,S}$ with $\dim \bar{Q}_d \geq 2$.*

Assuming the validity of this lemma, we complete the proof of Theorem 4.1. We only show (2) in this theorem; again (1) can be shown in a similar way. By this lemma, we can take a $\mathcal{C}$-minimal metric d in $\mathcal{D}_{\mu,S}$ with $\dim \bar{Q}_d \geq 2$. Take a cyclically tight extension d' of d such that d' contains $\beta\gamma_n$ as a submetric for sufficiently large n. Since d' is $\mathcal{C}$-minimal (Lemma 4.5 (2)), d' is decomposed, in modulo $\mathcal{L}$, into a convex combination of $\mathcal{C}$-minimal metrics $d_1, d_2, \dots, d_m$ such that each d_i is an extreme point of $\mathcal{D}'_{\mu,V}/\mathcal{L}$. Some $d_i + \mathcal{L}$ has no $1/k$-integral point, as in the metric case.

Let us start the proof. For $p \in P_\mu$, let X_p be the set of elements $s \in S$ with $p(s^c) + p(s^r) = 0$. Our argument crucially relies on the following claim:

(4.3) For $p \in P_\mu$, there is a minimal metric $d \in \mathcal{D}_{\mu,S}$ such that $p \in P_d$, and

$$\begin{array}{ll} p(s^c) + p(t^r) = d(st) & \text{if } s^c t^r \in EK_\mu(p), \\ p(s^c) + p(t^r) > d(st) & \text{otherwise,} \end{array} \quad (s, t \in S \setminus X_p).$$

Proof. Replacing p by $p + \alpha(\mathbf{1}, -\mathbf{1})$ for some α if necessary, we may assume that p is nonnegative. Let d be the distance on S defined by

$$d(st) = p(s^c) + p(t^r) \quad (s, t \in S).$$

Then d is a metric, more precisely, d is realized by a subdivision of a star. We try to decrease $d(st)$ for $s, t \in S \setminus X_p$ with $s^c t^r \notin EK(p)$ with keeping the triangle inequality. Since one of $p(s^c), p(s^r)$ and one of $p(t^c), p(t^r)$ are positive, there is no $u \in S \setminus \{s, t\}$ such that $d(us) + d(st) = d(ut)$ or $d(st) + d(tu) = d(su)$. Let $d(st) \leftarrow d(st) - \epsilon$ for small $\epsilon > 0$ and $s, t \in S \setminus X_p$ with $s^c t^r \notin EK(p)$; we remark $d(st) > \mu(st) \geq 0$. Then d does not violate the triangle inequality. So d is not minimal, and we can take a minimal metric $d' \in \mathcal{D}_{\mu,S}$ with $d' \leq d$. □

Proof of Lemma 4.6 (1). By (2.7) (2), we can take a point $p \in T_\mu$ such that $K_\mu(p)$ has at least k components having no $u \in S^{cr}$ with $p(u) = 0$. Take a minimal metric d in (4.3). Let $U := S \setminus X_p$. Consider the restrictions μ_U of μ to U and p_U of p to U^{cr}. Then p_U belongs to T_{μ_U} and $K_{\mu_U}(p_U)$ has k components. By (2.7) (2) we have $\dim T_{\mu_U} \geq k$. One can easily see that T_{μ_U} is the surjective image of the projection of T_μ; necessarily $\dim T_\mu \geq \dim T_{\mu_U} \geq k$.

Proof of Lemma 4.6 (2). Suppose that $\dim \bar{Q}^{slim}_\mu = k$ for $k \geq 2$. We try to find a triple (U, d, p) of $U \subseteq S$, a metric d on U, and $p \in Q_d$ such that d is $\mathcal{C}$-minimal in $\mathcal{D}_{\mu_U,U}$ and $K_d(p)$ has at least 3 components, which implies Lemma 4.6 (2) by (2.7) (2') and the following claim.

(4.4) For $U \subseteq S$, let d be a $\mathcal{C}$-minimal metric in $\mathcal{D}_{\mu_U,U}$. Then there is a $\mathcal{C}$-minimal metric d^* in $\mathcal{D}_{\mu,S}$ with $\dim \bar{Q}_{d^*} \geq \dim \bar{Q}_d$.

Proof. Extend d to d' in $\mathcal{D}_{\mu,S}$ with ${d'}_U = d$. Then d' may not be $\mathcal{C}$-minimal in $\mathcal{D}_{\mu,S}$. We can take a $\mathcal{C}$-minimal metric d^* in $\mathcal{D}_{\mu,S}$ such that $d^*(C) \leq d(C)$ for all cycles C in S. Since d is $\mathcal{C}$-minimal in $\mathcal{D}_{\mu_U,U}$, we have $d \equiv {d^*}_U \mod \mathcal{L}$. Then Q_d is a translation of $Q_{{d^*}_U}$, and hence $\dim \bar{Q}_d = \dim \bar{Q}_{{d^*}_U}$. Since every point $p \in Q_{{d^*}_U}$ can be extended to a point in Q_{d^*}, we have $\dim \bar{Q}_{d^*} \geq \dim \bar{Q}_d$. $\square$

We can take $p \in Q^{slim}_\mu$ such that $p/\sim$ belongs to the interior of k-dimensional face in $\bar{Q}^{slim}_\mu$ for $k \geq 2$. There are three cases:

(i) $K_\mu(p)$ has at least 3 components and $X_p = \emptyset$.

(ii) $K_\mu(p)$ has at least 3 components and $X_p \neq \emptyset$ (p belongs to Q_{μ,X_p}).

(iii) $K_\mu(p)$ has at least 4 components one of which is a (complete bipartite) component of nodes ${X_p}^c \cup {X_p}^r$ (p belongs to the interior of Q^{deg}_{μ,X_p}).

In the following, for a distance g on S and a point $p \in \mathbf{R}^{S^{cr}}$, we denote by $H_g(p)$ the directed graph on S with $EH_g(p) := \{st \mid s^c t^r \in EK_g(p)\}$ (possibly including loops).

We first consider case (i). Suppose that $H_\mu(p)$ is strongly connected. Since $X_p = \emptyset$, by (4.3) we can take a minimal metric $d \in \mathcal{D}_{\mu,S}$ and $K_\mu(p) = K_d(p)$. So $K_d(p)$ also has at least 3 components; $\dim \bar{Q}_d \geq 2$. By definition and construction, $EH_\mu(p) = EH_d(p) \subseteq EH_{\mu,d}$. Therefore $H_{\mu,d}$ is also strongly connected, which immediately implies the $\mathcal{C}$-minimality of d by Lemma 4.3 (2). Thus (S, d, p) is a required triple.

Suppose that $H_\mu(p)$ is not strongly connected. Take a chordless cycle $(x_1, x_2, \ldots, x_m)$ (without repeated nodes). Since $X_p = \emptyset$, equivalently, $H_\mu(p)$ has no loop, the cycle length m is at least two. Suppose $m \geq 3$. Then let $U \leftarrow \{x_1, x_2, \ldots, x_m\}$, and $p \leftarrow p_U$ (the restriction of p to U^{cr}). Then $H_{\mu_U}(p)$ is a cycle of length m, and $K_{\mu_U}(p)$ is a matching of size m (has m components). According to (4.3), we can take a minimal metric $d \in \mathcal{D}_{\mu_U,U}$ with $p \in Q_d$ and $K_{\mu_U}(p) = K_d(p)$. Thus $H_{\mu_U,d}$ is strongly connected, and (U, d, p) is a required triple.

Suppose that there is no simple chordless cycle of length at least 3 in $H_\mu(p)$. Since each node has both entering and leaving edges, we can take from $H_\mu(p)$ two disjoint 2-cycles (x_1, x_2), (y_1, y_2) without edges from $\{y_1, y_2\}$ to $\{x_1, x_2\}$. Let $U \leftarrow \{x_1, x_2, y_1, y_2\}$, and $p \leftarrow p_U$. Again we can take a minimal metric $d \in \mathcal{D}_{\mu_U,U}$ with $K_{\mu_U}(p) = K_d(p)$. For small positive ϵ, decrease p by ϵ on $\{y_1, y_2\}^c$ and increase p by ϵ on $\{y_1, y_2\}^r$. Then $K_d(p)$ has four components consisting of four disjoint edges $x_1^c x_2^r, x_2^c x_1^r, y_1^c y_2^r, y_2^c y_1^r$. Suppose that d is not $\mathcal{C}$-minimal (otherwise (U, d, p) is a required triple). So we may assume that $H_{\mu_U,d}$ has an extremal edge from $\{x_1, x_2\}$ to $\{y_1, y_2\}$, and has no edge from $\{y_1, y_2\}$ to $\{x_1, x_2\}$. Then we can construct a $\mathcal{C}$-minimal metric from d by adding

$$\begin{array}{|c|cccc|}
\hline
 & x_1 & x_2 & y_1 & y_2 \\
\hline
x_1 & 0 & 0 & \epsilon & \epsilon \\
x_2 & 0 & 0 & \epsilon & \epsilon \\
y_1 & -\epsilon & -\epsilon & 0 & 0 \\
y_2 & -\epsilon & -\epsilon & 0 & 0 \\
\hline
\end{array} \in \mathcal{L}$$

for $\epsilon > 0$ and decreasing some of $d(x_i y_j)$. For keeping $p \in Q_d$, decrease p by ϵ on $\{y_1, y_2\}^c$ and increase p by ϵ on $\{y_1, y_2\}^r$. Then the resulting $K_d(p)$ is a matching consisting of the four edges. Thus (U, d, p) is a required triple.

Next we consider case (iii). Let $U \leftarrow S \setminus X_p$ and let $p \leftarrow p_U$. Then $K_{\mu_U}(p)$ has no isolated node. Obviously, $K_{\mu_U}(p)$ has at least 3 connected components, and $X_p = \emptyset$. Thus $p \in Q^{slim}_{\mu_U}$. Therefore, the situation reduces to case (i).

Finally we consider case (ii). We use the same projection idea. Take $x \in X_p$. Let $A^c \subseteq S^c \setminus {X_p}^c$ and $B^r \subseteq S^r \setminus {X_p}^r$ be the sets of nodes u covered only by x^r and x^c, respectively. Suppose $A^c \cup B^r = \emptyset$. Then let $U \leftarrow S \setminus x$, and let $p \leftarrow p_U$. Then X_p decreases, $p \in Q^{slim}_{\mu_U}$, and $K_{\mu_U}(p)$ has at least 3 connected components. Suppose that both A^c and B^r are nonempty. For a small positive ϵ, decrease p by ϵ on $A^c \cup B^r$ and increase p by ϵ on x^c, x^r. Then X_p decreases, $p \in Q^{slim}_{\mu}$, and $K_\mu(p)$ has at least 5 connected components. Suppose that one of A^c and B^r, say B^r, is empty. Then x^c is necessarily incident to $S^r \setminus x^r$; otherwise p is a proper fat relative to $\{x\}$, a contradiction. For a small positive ϵ, decrease p by ϵ on A^c and increase p by ϵ on x^r. Again $p \in Q^{slim}_{\mu}$ (x^c is still covered), and $K_\mu(p)$ has at least 4 connected components; Repeat it until $X_p = \emptyset$. After that, the situation reduces to case (i).

Acknowledgements

We thank the referee for careful reading and helpful comments. The first author is supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan. The second author is supported by Nanzan University Pache Research Subsidy I-A-2 for the 2008 academic year.

References

[1] M. Develin and B. Sturmfels, Tropical convexity, *Documenta Mathematica* **9** (2004), 1–27.

[2] A. W. M. Dress, Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: a note on combinatorial properties of metric spaces, *Advances in Mathematics* **53** (1984), 321–402.

[3] A. Frank, On connectivity properties of Eulerian digraphs, In: *Graph theory in memory of G. A. Dirac* (Sandbjerg, 1985; L. Døvling Andersen, I. Tafteberg Jakobsen, C. Thomassen, B. Toft, P.D. Vestergaad, eds.), North-Holland, Amsterdam, 1989, pp. 179–194.

[4] H. Hirai, Characterization of the distance between subtrees of a tree by the associated tight span, *Annals of Combinatorics* **10** (2006), 111–128.

[5] H. Hirai, Tight spans of distances and the dual fractionality of undirected multiflow problems, *Journal of Combinatorial Theory, Series B* **99** (2009), 843–868.

[6] H. Hirai, Folder complexes and multiflow combinatorial dualities, RIMS-Preprint 1675, (2009).

[7] H. Hirai, The maximum multiflow problems with bounded fractionality, RIMS-Preprint 1682, (2009).

[8] H. Hirai and S. Koichi, On tight spans and tropical polytopes for directed distances, preprint 2010, `arXiv:1004.0415`.

[9] T. Ibaraki, A. V. Karzanov, H. Nagamochi, A fast algorithm for finding a maximum free multiflow in an inner Eulerian network and some generalizations, *Combinatorica* **18** (1998), 61–83.

[10] J. R. Isbell, Six theorems about injective metric spaces, *Commentarii Mathematici Helvetici* **39** (1964), 65–76.

[11] A. V. Karzanov, Polyhedra related to undirected multicommodity flows, *Linear Algebra and Its Applications* **114/115** (1989), 293–328.

[12] A. V. Karzanov, Minimum 0-extensions of graph metrics, *European Journal of Combinatorics* **19** (1998), 71–101.

[13] A. V. Karzanov, Metrics with finite sets of primitive extensions, *Annals of Combinatorics* **2** (1998), 211–241.

[14] M. V. Lomonosov, unpublished manuscript, 1978.

[15] A. Schrijver, *Theory of Linear and Integer Programming*, Wiley, Chichester, 1986.